

**The Dichotic Digits difference Test (DDdT): Development,
Normative Data and Retest Reliability Studies**

Journal:	<i>Journal of the American Academy of Audiology</i>
Manuscript ID:	Draft
Manuscript Type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Cameron, Sharon; National Acoustic Laboratories, Assessment Glyde, Helen; National Acoustic Laboratories, Assessment Dillon, Harvey; National Acoustic Laboratories, Assessment Whitfield, Jessica; National Acoustic Laboratories, Assessment Seymour, John; National Acoustic Laboratories, Assessment
Keywords:	Auditory processing disorder, Diagnostic techniques, Hearing science

SCHOLARONE™
Manuscripts

view

Submission date: 10 August 2015

The Dichotic Digits difference Test (DDdT): Development, Normative Data and Retest
Reliability Studies

Sharon Cameron ^a
Helen Glyde ^a
Harvey Dillon ^a
Jessica Whitfield ^a
John Seymour ^a

^aNational Acoustic Laboratories

Parts of the research described in this paper were presented at the XXXII World Congress of Audiology, Brisbane, Australia, May 2014.

Acknowledgments. The authors would like to thank Mark Seeto for statistical advice. Thanks are also due to the Catholic Schools Office and the various primary schools who took part in this research. The participation of the children and their families are also appreciated. This research is funded by the Australian Government through the Department of Health.

Corresponding Author:

Sharon Cameron, PhD
Senior Research Scientist
National Acoustic Laboratories
Australian Hearing Hub
16 University Avenue
Macquarie University NSW 2109
Australia
Phone: +61 2 9412 6851
Fax: +61 2 9412 6769
e-mail: Sharon.Cameron@nal.gov.au

Abstract

Background: The dichotic digits test is one of the most widely used assessment tools for central auditory processing disorder (CAPD). However, questions remain concerning the impact of cognitive factors on test results.

Purpose: To develop the Dichotic Digits difference Test (DDdT), an assessment tool which could differentiate children with cognitive deficits from children with genuine dichotic deficits based on differential test results. The DDdT consists of four subtests: dichotic free recall (FR), dichotic directed left (DLE) and right ear (DRE), and diotic. Scores for six conditions are calculated (Free Recall LE, RE and total, as well as the DLE, DRE and diotic). Scores for four difference measures are also calculated: Dichotic advantage, Right Ear Advantage (REA) FR; REA Directed and Attention Advantage.

Research Design: Experiment 1 involved development of the DDdT, including error rate analysis. Experiment 2 involved collection of normative and test retest reliability data.

Study Sample: Twenty adults (25, 10 (yrs, mths) to 50, 7, mean 36, 4) took part in the development study; 62 normal-hearing, non-clinical, primary-school children (7, 1 (yrs, mths) to 11, 11, mean 9, 4) and ten adults (25, 0 (yrs, mths) to 51, 6, mean 34, 10) took part in the normative and retest reliability study.

Data Collection and Analysis: In Experiment 1 error rate analysis was conducted on the 36 digit pair combinations of the DDdT. Normative data collected in Experiment 2 was arcsine transformed to achieve a distribution that was closer to a normal distribution and z-scores calculated. Pearson product moment correlations were utilized to determine the strength of relationships between DDdT conditions.

Results: The development study revealed no significant differences in the adult population between test and retest on any DDdT condition. Error rates on 36 digit pairs ranged from 1.5% to 16.7%. The most and least error-prone digits were removed prior to commencement

1
2
3 of the normative data study, leaving unique 25 digit pairs. Average z-scores calculated from
4
5 the arcsine-transformed data collected from the 62 children who took part in the normative
6
7 data study revealed that free recall dichotic processing (LE, RE and total) was highly
8
9 correlated with diotic processing (r ranging from 0.5 to 0.6; $p < 0.0001$). Significant
10
11 improvements in performance on retest occurred for the FR LE, RE and total conditions (p
12
13 ranging from 0.05 to 0.0004), the conditions that would be expected to improve with practice
14
15 if the participant's response strategies are better the second time around.
16
17

18 **Conclusions:** The addition in the DDdT of a diotic control task - that shares many response
19
20 demands with the usual dichotic tasks - opens up the possibility of differentiating children
21
22 who perform badly because of poor dichotic processing from those who perform poorly
23
24 because of poor attention, memory, or other cognitive abilities. The high correlation between
25
26 dichotic and diotic performance suggests that factors other than dichotic performance play a
27
28 substantial role in a child's ability to perform a dichotic listening task. This hypothesis will be
29
30 investigated further in the cognitive correlation study presented as paper 2.
31
32
33
34
35
36
37

38 **Key Words:** Dichotic, diotic, free recall, directed; impulsivity; sustained attention; central
39
40 auditory processing disorder
41
42
43
44
45
46

47 **Abbreviations:** CAPD = Central Auditory Processing Disorder; DDdT = Dichotic Digits
48
49 difference Test; DLE = directed left ear; DRE = directed right ear; FR = free recall; NAL =
50
51 National Acoustic Laboratories; REA = right ear advantage; RMS = root mean square.
52
53
54
55
56
57
58
59
60

INTRODUCTION

Although the make-up of central auditory processing test batteries vary widely from clinic to clinic, one of the most common types of tests used are dichotic listening tests (Martin et al, 2007; Schmithorst et al, 2013; Weihing and Atcherson, 2014). Both the American Academy of Audiology (AAA, 2010) and the American Speech and Hearing Association (ASHA, 2005) recommend the inclusion of a dichotic listening task - either dichotic digits, words, or sentences - in the assessment of central auditory processing disorder (CAPD).

Dichotic processing refers to the simultaneous presentation through headphones of different acoustic signals to the left and right ears (Musiek and Weihing, 2011). It is different to diotic processing, in which the same signal is presented to both ears. Clinically, dichotic listening is evaluated using either a free recall task (whereby the patient attends to both ears and repeats back what is heard) or a directed task, whereby the patient is asked to attend to only one ear (Musiek and Weihing, 2011). The free recall task is defined as a measure of binaural integration whereas the directed task is a measure of binaural separation. For both tasks, results are typically calculated separately as a percentage correct for the left and right ear.

From an anatomical perspective, Musiek and Weihing (2011) explain that the right ear signal will travel directly to the left hemisphere of the cortex via the dominant contralateral pathways, whereas the left ear signal travels to the right hemisphere and then crosses to the left hemisphere via the corpus callosum for processing and verbal response. Thus effective interhemispheric transfer between the two cerebral hemispheres is an important component of dichotic processing. Speech-related stimuli presented to the right ear are typically recalled with greater accuracy than that presented to the left ear (Kimura, 2011; Schmithorst et al,

1
2
3 2013). This right ear advantage (REA) is said to be a consequence of preferential conduction
4
5 of right-ear messages directly to the typically language-dominant left hemisphere via the
6
7 stronger contralateral pathways (Hiscock and Kinsbourne, 2011; Kimura, 2011). However,
8
9 not all left-hemisphere dominant individuals report a REA (Schmithorst et al 2013) and the
10
11 magnitude of REA has been shown to decrease in older compared with younger children
12
13 (Moncrief, 2011).
14

15
16 According to Musiek and Weihing (2011) a significant left ear deficit (that is, a relatively
17
18 *increased* right ear advantage) on dichotic processing tasks is associated with a breakdown in
19
20 the interhemispheric transfer function at the level of the corpus callosum. Further, Bellis
21
22 (2003) suggests that binaural integration and separation processes are reported to be critical
23
24 to everyday listening and deficits in these areas, as measured by free recall and directed
25
26 dichotic tests, may be expressed behaviourally as difficulty hearing in background noise or
27
28 when more than one person is talking at the same time. Given the reported potential
29
30 behavioural consequences of dichotic deficits, assessment certainly seems warranted.
31
32
33

34 However, DeBonis (2015) notes that whereas dichotic tests have been included in
35
36 auditory processing batteries for over four decades, research has now established that
37
38 performance on dichotic listening tasks is greatly influence by allocation of attentional
39
40 resources. In a review of clinical and experiment evidence concerning dichotic listening
41
42 studies of hemispheric asymmetry, Westerhausen and Hugdahl (2008), concluded that the
43
44 corpus callosum is not only a channel for the automatic exchange of information between the
45
46 cerebral hemispheres, but also allows for dynamic and flexible interaction that supports both
47
48 bottom-up and top-down processing of auditory stimuli. That is, dichotic processing involves
49
50 a built-in, automated (or stimulus-driven) component which favours the processing of right
51
52 ear input, as well as a cognitive (instruction-driven) component that allows for modulation of
53
54 the laterality effect. To this end, Hiscock and Kinsbourne (2011) state that REA effects
55
56
57
58
59
60

cannot be explained on the basis of hypothetical conduction properties of specific auditory pathways, but instead in terms of dynamic processes, including those in which individuals deliberately direct their attention in different ways. As such, the authors conclude that the role of attention is essential for interpreting the process of REA.

In a discussion of resource allocation theory, Bellis (2014) points out, that individuals with higher-order attention, cognitive or related deficits may exhibit decreased performance in tests of fundamental sensory processing due to the extra allocation of effort necessary to attend to, comprehend or remember the stimulus. To this end, Ahmmed et al (2014) reported that in a sample of 110 children with suspected CAPD, performance on a dichotic free recall word test correlated significantly with performance on tests of forward and reverse digit span ($r = 0.34$, $p < 0.01$), and auditory attention cued ($r = 0.30$, $p < 0.01$) and non-cued ($r = 0.23$, $p < 0.05$), but not with non-verbal IQ ($r = -0.03$). Similarly, in a sample of 36 children with suspected CAPD, Maerlender (2010) found a significant correlation between dichotic free recall left ear and digit span forward ($r = 0.68$, $p < 0.001$) and reversed ($r = 0.45$, $p < 0.01$), as well as free recall right ear and digit span forward ($r = 0.38$, $p = 0.02$) and reversed ($r = 0.40$, $p = 0.01$). The author concluded that verbal memory may be a pervasive deficit in children with CAPD, and the value and role of psychometric testing in children suspected of this disorder are supported. It seems more appropriate, however, to conclude that verbal memory deficit may be pervasive in children for whom an assessment for CAPD is sought.

In an investigation of links between auditory processing test results, functional deficits and cognitive abilities in a sample of 50 control and 105 children referred for auditory processing assessment, Tomlin et al (2015) found that both left and right ear free recall dichotic digit test scores correlated significantly ($p \leq 0.01$) with forward and reverse digit span and non-verbal IQ, as well as with attention ($p \leq 0.01$ for left ear and $p \leq 0.05$ for right ear). Multiple regression analysis found that auditory working memory was a significant

1
2
3 predictor of dichotic left and right ear performance ($\beta = 0.42$, $p < 0.001$; and $\beta = 0.24$, $p <$
4
5 0.01 respectively) and that non-verbal IQ was a significant predictor of left ear free recall
6
7 performance ($\beta = 0.58$, $p < 0.01$). Tomlin et al (2015) concluded that interpretation of
8
9 auditory processing tests requires consideration of how cognitive ability may have impacted
10
11 on both test results and the functional difficulties experienced by the child.

12
13
14 The aim of the present study was to develop a test that could be used clinically to
15
16 identify children with true dichotic processing deficits from those with other deficits which
17
18 may account for poor dichotic free recall and/or directed performance. The Dichotic Digits
19
20 difference Test (DDdT; Cameron et al, 2013) utilizes a diotic control condition and
21
22 measurement of difference scores to alert clinicians to a potential impact of higher order
23
24 deficits on test performance. The DDdT scoring is highly automated as described in the
25
26 Methods section to follow. This paper – part one of two on the evaluation of the DDdT – will
27
28 report on error analysis of digit pairs during the development phase of the study, as well as
29
30 provide normative and retest reliability data in a pediatric and adult population. Correlations
31
32 between DDdT condition scores are also investigated. The second paper will investigate
33
34 correlations between DDdT conditions and measures of cognitive ability (attention, memory
35
36 and intelligence) as well as measures of real life listening ability.
37
38
39
40
41
42
43
44

45 **EXPERIMENT 1 – ADULT DEVELOPMENT STUDY**

46 **DDdT Software Development**

47
48
49 The DDdT graphical user interface and signal processing application were developed in
50
51 MATLAB programming language (MathWorks, 2013). An image of the user interface is shown
52
53 in Figure 1. The monosyllabic digits 1, 2, 3, 4, 5, 6, 8, 9 and 10 were recorded in an anechoic
54
55 chamber at the National Acoustic Laboratories (NAL). The stimuli were voiced by an adult
56
57
58
59
60

male with an Australian English accent. Each number was spoken twice in a normal clear voice. The better of the two recordings for each digit, as judged subjectively by two of the authors, were then edited to remove extraneous silence from the beginning and end of each file. This resulted in files with an average length of 631 ms. Each digit was level normalized to a mean RMS level of -25 dB re full digital level to ensure no clipping occurred during playback. To calibrate the software a 1 kHz calibration tone with a mean RMS level of -15 dB re full digital level was activated from the playback screen and the volume of the laptop was adjusted until the electrical level of the calibration signal was 18 mV. This resulted in an RMS presentation level of 50 dB (HL) of the digits when presented through Sennheiser HD215 headphones.




Figure 1. DDdT graphical user interface (research version).

Method

Approval for the study was granted from the Australian Hearing Human Research Ethics Committee.

Participants

Twenty adults with normal hearing, defined as equal to or better than 20 dB HL at all octave frequencies between 250 Hz and 8000 Hz, were recruited to participate in the study. Participants were aged 25, 10 (yrs, mths) to 50, 7 (mean 36, 4) and had no reported history of learning or attention disorders. There were five males and fifteen females. All participants spoke English as their first language. Thirteen participants were retested between 14 days and 31 days after their initial appointment to investigate retest reliability.

Procedure

Testing was conducted in a sound-attenuated booth at the National Acoustic Laboratories and took approximately 30 minutes to complete. Pure tone audiometric screening was performed using a Maico MA 53 clinical audiometer with circumaural Sennheiser HDA 200 audiometric headphones.

The DDdT was administered with the use of a laptop computer and Sennheiser HD215 headphones. Sound levels were calibrated prior to each appointment, using the procedure described above. The four subtests of the DDdT (free recall (FR), directed left ear (DLE), directed right ear (DRE) and diotic) were counterbalanced between participants. Five practice trials and twenty scored trials were presented in each subtest and the participants' task varied depending on the subtest. Each trial consisted of four randomly selected digits presented as temporally overlapped pairs presented. The four subtests were presented, as described below, resulting in six conditions, which are scored as percent correct:

- 1
2
3 a) Dichotic Free Recall (FR LE, FR RE and FR Total): a set of two numbers were presented
4 to the left ear (e.g. 8, 6) at the same time as two *different* numbers were presented to the
5 right ear (e.g. 1, 3). The digits presented first to each ear overlapped in time, as did the
6 digits presented second to each ear. Participants were asked to repeat back all four
7 numbers in any order (e.g. 6, 8, 1, 3). The LE and RE were scored separately. The
8 average of the LE and RE scores – referred to as FR Total - was also calculated.
9
10
11
12
13
14
15
16
17 b) Dichotic directed LE (DLE): Presentation of stimuli was as per Dichotic FR described
18 above. However participants were asked to repeat back only the digit pair heard in the
19 left ear (e.g. 8, 6), in any order.
20
21
22
23
24
25 c) Dichotic directed RE (DRE): As for DLE, however participants were asked to repeat
26 back only the digit pair heard in the right ear (e.g. 1, 3), in any order.
27
28
29
30
31 d) Diotic: A series of four numbers were presented to *both* ears (e.g. 8, 1, 6, 3). Two digits
32 overlapped exactly in time (e.g. 8, 1), followed by the other two overlapping digits (6, 3).
33 Participants were asked to repeat back all four digits in any order.
34
35
36
37
38

39 Results

40
41 Statistical Analysis was performed using Statistica Version 10 and Excel 2013.
42
43
44

45 *Analysis for effects of order*

46
47 A mixed-effects logistics regression model was fitted to the data to determine the effects
48 of test order on condition score. The results indicated that, when averaged across the various
49 DDdT conditions, test scores increased with increased position in test order, with position 3
50 and 4 significantly higher than position 1 ($p = 0.3$ and 0.01 respectively). When merged
51 across test orders, Friedman ANOVA revealed that the mean score in the diotic condition
52
53
54
55
56
57
58
59
60

(96.1%) was significantly lower ($p = 0.02$) than in the dichotic FR Total condition (97.9%).

There were no other significant differences between any other combination of mean scores (FR LE 97.3%; FR RE 98.6%; DLE 98.1%; DRE 98.9%). Inter-subject variation ranged from 2.1 standard deviations (SD) on DRE and FR RE to 3.9 SD on the dichotic condition.

Error Rates Analysis

To determine whether all pair combinations were equally difficult error rates were analysed from the DDdT free recall and diotic results. Figure 2 shows the percentage of presentations in which an error was made for each of the 36 pair combinations, collapsed across conditions. The average error rate was 6.15%, but the error rate for individual pairs varied from 1.5% to 16.7%.




Figure 2. Percent of incorrect response for each digit pair combination for the 20 adults in the development study.

Test-Retest Reliability Analysis

Repeated measures ANOVA revealed that there were no significant differences between test and retest on any of the DDdT test scores (p ranging from 0.08 to 0.83). Any small differences in performance by the 13 adults who took part in the retest reliability study were in the direction of improvement at retest. The largest improvement on retest occurred in the diotic condition, where performance improved on average from 96.7% correct at test to 98.1% correct at retest.

Modifications to Test Based on Results

Given that the dichotic free recall scores are higher than the diotic score it appears that dichotic presentation facilitates the understanding of temporally overlapped digits. Given this, it was determined that further research with the DDdT to ascertain its value as a clinical tool for differentiating memory or attention deficits from dichotic deficits was warranted. However, to ensure that each time the test was used with a different random set of digit pairs, the level of difficulty remained constant it was necessary to remove digit pairs which were likely to be markedly easier or more difficult than the remaining pairs. To this end the five most error prone and six least error prone number combinations (1/2, 1/9, 2/4, 2/6, 3/8, 3/9, 4/10, 5/6, 6/8, 8/9, & 9/10) were removed. This left 25 permitted digit pairs. A new random sampling rule was implemented in which each test condition would contain all permitted digit pairs appearing twice within the 25 trials (5 practice and 20 scored). Given the difficulty level of the permitted pairs is approximately equal, the pairs are allowed to appear in either the practice or scored position. Once these changes were implemented an adult normative data study was completed. Further, based on the analysis for effects of order and condition, DDdT conditions were presented in a fixed order (FR, DLE, DRE and diotic) in the normative data study described below.

EXPERIMENT 2 – NORMATIVE DATA AND RETEST RELIABILITY STUDY

Method

Approval for the study was granted from the Australian Hearing Human Research Ethics Committee and the Catholic Schools Office, Diocese of Broken Bay.

Participants

A total of 72 participants took part in the study. There were ten adults aged 25, 0 (yrs, mths) to 51, 6 (mean 34, 10), of which three were male and seven were female. Five of the adult participants had previously taken part in the development study described above. For these participants a minimum of twelve months had elapsed between the two studies to minimize the potential impact of previous exposure to the test procedure on results.

There were 62 children aged 7, 1 (yrs, mths) to 11, 11 (mean 9, 4) who took part in the study, of which 27 were male and 35 were female. The children were recruited from a Catholic primary school in New South Wales, Australia, with an average Index of Community Socio-Educational Advantage value similar to the national average. All participants spoke English as their first language, had no reported history of diagnosed or suspected learning or attention disorders and had normal hearing, defined as equal to or better than 20 dB HL at all octave frequencies from 250 Hz to 8000 Hz. Thirty children were re-assessed on the DDdT two months after initial testing to determine retest reliability.

Procedure

Adults were assessed at NAL as described in Experiment 1 with the exception that the DDdT conditions were no longer counterbalanced, but presented in the following order: free recall (FR), directed left ear (DLE), directed right ear (DRE) and diotic. Children were assessed in a quiet room on the school grounds. Sound levels in the school testing rooms were measured between 45-50 dBA using a Q1362 digital sound level meter. Pure tone audiometric screening was performed using an Interacoustics Audio Traveller A222 portable audiometer with Telephonics TDH 39P audiometric headphones in H7A Peltor cups. The DDdT was presented to the adults and children as described in Experiment 1.

Results

Statistical Analysis was performed using Statistica Version 10. As the data were percent correct scores, often close to or at 100%, data were arcsine transformed prior to analysis to more closely approximate a normal distribution.

Gender and Age

The effect of gender on performance was examined in the pediatric data. There was no significant difference between males and females on any of the six DDdT conditions, ranging from $F_{(1,60)} = 1.060, p = 0.307$ for the Directed LE condition to $F_{(1,60)} = 3.291, p = 0.074$.

The mean percentage scores for all participants as a function of age group are provided in Table 1. Transformation of the data and analysis of the effects of age are discussed in the following section.

Table 1. Average percent correct scores on the six DDdT conditions as a function of age.

	Age					
	7	8	9	10	11	Adult
FR LE %	79.5	73.2	79.6	88.8	90.6	99.8
FR RE %	88.9	85.0	86.3	91.8	91.7	99.8
FR Total %	84.4	79.1	82.9	90.3	91.1	99.8
DLE %	81.3	86.1	86.5	91.6	94.4	99.8
DRE %	91.6	87.1	91.9	95.0	96.7	99.3
Diotic %	72.1	72.0	77.3	83.2	87.5	96.0

Arcsine Transformation, Regression and Creation of Z-Scores

The normative data collected from 62 children and 10 adults were used to create equations that allow the expression of individual scores in age-corrected population standard deviation units, that is, z-scores. The method described in Tomlin et al (2014) was used. In brief, this comprised arcsine transformation of the scores, non-linear regression of the transformed scores against age, and calculation of the root mean square value of the residual deviations from the fitted functions.

The percentage scores for six DDdT conditions - FR LE, FR RE, FR Total (average of LE and RE); DLE, DRE and diotic - were arcsine transformed using equation 1:

$$T = \text{sine}^{-1}(\sqrt{p}) \dots 1,$$

where p is the proportion of digits correct and T is the transformed score. These transformed scores were then regressed against age using the exponential formula shown in equation 2:

$$T' = a - b.e^{(-age/c)} \dots 2,$$

where T' is the estimated value of T and a , b , and c are the coefficients that determine each curve.

The proportion of variance in transformed scores accounted for by the regression varied from 26% for the DRE condition up to 60% for the FR Total condition. The higher proportion of variance accounted for in the case of the FR Total scores likely reflects the smaller error variance in the total scores, due to their being based on twice as many items as for the individual ear scores.

A preliminary fitting indicated that the value of c varied from 4.5 years for the DRE condition up to 17.3 years for the FR RE condition. However, because there were no participants with ages between 13 and 25 years, the value of c could be varied widely, along with a compensating variation in the values of a and b , without markedly affecting the fit to the data. Consequently, all six curves were re-fit with c fixed at its median value (7.42 years) found across the conditions in the preliminary fitting. The resulting values of a and b are shown in Table 2.



Because the scores for the adults were close to ceiling, the residual scores (transformed score minus predicted transformed score) were much smaller for the adults than for the children. Because of this difference, and because of small number of adult subjects, further analysis of the residual scores was carried out for the child data only. The squared residual errors were linearly regressed against age. No significant correlations emerged, and the largest correlation was only -0.12. Consequently, the residual error was characterized by its standard deviation (σ), independent of age. These standard deviations are also shown in Table 2. Calculation of z-scores was then carried out in the usual manner, but using transformed scores and predicted scores, rather than raw scores, as shown in Equation 3:

$$z = (T - T') / \sigma \dots 3,$$



Table 2. Regression coefficients for the exponential formula shown in Equation 2 ($T' = a - b.e^{(-age/c)}$). Also shown is σ , the standard deviation of the residuals (transformed scores minus predicted transformed scores).

Condition	<i>a</i>	<i>b</i>	σ
FR LE	1.572	1.410	0.160
FR RE	1.542	0.972	0.148
FR Total	1.546	1.201	0.120
DLE	1.581	1.180	0.149
DRE	1.565	0.786	0.161
Diotic	1.440	1.201	0.116

The regression curves (Equation 2) can be inverse transformed back to percentage values (i.e. percent correct) using the inverse of Equation 1. These are shown in Figure 3 (a to f), along with the original data, also in percentage values. Also shown in these graphs are the percentage values that correspond to the transformed values that are two standard deviations below the age-dependent mean values. These are shown only up to age 12 as they are based on the residual values only up to this age.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60






Figure 3. Scatterplots of the untransformed DDdT data and the regression curve (solid line) transformed back to percentage values (i.e. percent correct): (a) FR LE; (b) FR RE; (c) FR Total; (d) DLE; (e) DRE; (f) Diotic, based on the data from the 62 children and 10 adults in

1
2
3 the normative data study. The dashed line represents the percentage values that correspond to
4
5 the transformed values that are two standard deviations below the age-dependent mean
6
7 values.
8
9

10 11 12 *Normality*

13
14 The z-scores were examined to determine if they deviated significantly from normal
15
16 distributions. Based on the Shapiro-Wilks test of normality, there was no evidence of a
17
18 deviation from normality for five of the 6 conditions, with p values ranging from 0.27 to 0.81.
19
20 The exception was the DRE condition, for which the Shapiro-Wilks W-value was 0.95 and
21
22 the corresponding p-value was 0.03. The greatest deviation from normality was a greater
23
24 than expected number of z-scores in the range 1 to 1.5 standard deviations above the mean.
25
26 These were caused by a number of 9 to 12-year olds all achieving a score of 100% in this
27
28 condition.
29
30
31
32
33

34 35 *Difference Scores*

36
37 The data were used to derive the following difference scores, D, between pairs of
38
39 conditions.
40

- 41 1. Dichotic advantage: Dichotic FR Total score minus diotic score.
 - 42 2. Right ear advantage (Free Recall): Dichotic FR RE score minus FR LE score.
 - 43 3. Right ear advantage (Directed): Directed RE score minus Directed LE score.
 - 44 4. Attention advantage: Average of the Directed RE and LE scores minus FR Total score.
- 45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Difference Statistics Based on Transformed Data

For each difference score, all of which were computed based on the transformed data (i.e. using Equation 1), the difference score was linearly regressed against age, using just the child data, and the regression coefficients f and g , as defined in Equation 4 and as shown in Table 3, were found:

$$TD' = f + g \cdot \text{age} \quad \dots 4,$$

where TD' is the expected age-appropriate difference in transformed scores. Of the four difference scores, only the right ear advantage score in the free recall condition was significantly correlated with age ($r=-0.26$, $p=0.04$). The standard deviation, σ , of the residual values was computed, enabling z-scores to be calculated as shown in Equation 5, values for which are also shown in Table 3:

$$z = (TD - TD') / \sigma \quad \dots 5.$$

Table 3. Regression coefficients for the equation shown in Equation 4. Also shown is σ , the standard deviation of the residuals (difference scores minus predicted difference scores).

Difference score	f	g	σ
Dichotic advantage	0.247	-0.0155	0.101
REA – free recall	0.390	-0.0317	0.173
REA – directed	0.176	-0.0081	0.182
Attention advantage	-0.003	0.0106	0.121

Difference statistics based on original data

Untransformed difference scores (i.e. in percentage points) were also regressed against age, because of the complications in inverse transforming a difference of two arcsine-transformed scores. Equation 6 was fitted to the untransformed data for the children and adults combined:

$$D' = h + i \cdot \exp(-age/j) \quad \dots 6.$$


The regression coefficients are shown in the Table 4, along with the standard deviation of the residuals. In the case of the two right-ear advantage measures, the adult data showed a right-ear advantage so close to zero, the h parameters were set exactly to zero prior to fitting. In the case of attention advantage, there was no apparent dependence on age, so the i parameter was set to zero. In all cases, the standard deviation of the residuals were calculated using only the child data.

Table 4. Regression coefficients for the equation shown in Equation 6. Also shown is σ , the standard deviation of the residuals (difference scores minus predicted difference scores).


Difference score	h	i	j	σ
Dichotic advantage	3.10	61.12	3.50	7.60
Right ear advantage – free recall	0	176.20	2.75	9.45
Right ear advantage – directed	0	50.41	3.91	10.48
Attention advantage	4.43	0	-	7.45

Figure 4 (a to d) shows the means and ± 2 standard deviation (SD) limits for the difference measures as a function of age.


(a)



(b)



(c)



(d)




Figure 4. Scatterplots of performance on the DDdT difference measures (Dichotic Advantage, Free Recall REA, Directed REA and Attention Advantage) by the 62 children in the normative data study. The solid line indicates the mean score as a function of age, the dashed lines show the + and - 2 standard deviation limits.

Test-retest Reliability

Repeated measures ANOVAs were performed on the data from the 30 children who took part in the test retest reliability study. Means, SDs and p values calculated from the transformed z-scores for the six DDdT conditions and four advantage measures are provided in Table 5. The mean differences (retest score minus test score) are also provided, together with the intra-subject variability, which provides a measure of the spread around the change from test to retest.

It was found that dichotic free recall improved significantly with practice (FR LE [$F_{(1,29)} = 4.008, p = 0.055$]; FR RE [$F_{(1,29)} = 7.420, p = 0.011$]; and FR Total [$F_{(1,29)} = 15.957, p = 0.0004$]). The diotic condition also improved significantly with practice ($F_{(1,29)} = 11.212, p = 0.002$). However, the directed conditions did not improve significantly with practice (DLE [$F_{(1,29)} = 0.014, p = 0.906$] and DRE [$F_{(1,29)} = 0.011, p = 0.918$]). Right ear advantage did not improve for either the free recall ($F_{(1,29)} = 0.005, p = 0.945$), or directed ($F_{(1,29)} = 0.001, p = 0.976$) conditions. Attention advantage decreases somewhat ($F_{(1,29)} = 3.906, p = 0.058$), because free recall performance improved with practice but directed performance did not.

Pearson Product Moment correlations between DDdT test and retest performance are also provided in Table 5. The largest test-retest correlations were found for the FR Total ($r = 0.72, p = 0.00001$) and Diotic conditions ($r = 0.71, p = 0.00001$).

The second-last column in Table 5 shows the percentage of score variance that likely reflects random measurement error. These were computed as one minus half the square of the test-retest standard deviations (given that the expected value of the total score variance is 1.0 because these are standardized scores, and that both test and retest measurement error contribute to the test-retest standard deviations). The two measurement results with the smallest percentage of measurement error are the FR Total score and diotic score, as each is

based on 80 scored items, whereas the other base scores are based on only 40 scored items. The advantage measures generally have a high proportion of measurement error, which is not surprising given that they are based on the difference between two scores.

Table 5. Average test, retest and retest minus test z-scores and SDs for each DDdT condition and advantage measure for the 30 children who took part in the test-retest reliability study. P-values from repeated measures ANOVAs of test and retest z scores and error variance (%) are also provided. Pearson product-moment correlation coefficient (r) is shown, with significant test-retest correlations marked with an asterisk (* < 0.05; ** < 0.01).

Condition	<i>Test</i>	<i>Retest</i>	<i>Retest-Test</i>		<i>p-value</i>	<i>Error variance (%)</i>	<i>r</i>
	<i>Mean z-score</i>	<i>Mean z-score</i>	<i>Mean z-score</i>	<i>SD</i>			
FR LE	-0.08	0.30	0.39	1.06	0.055	56	0.36 *
FR RE	-0.24	0.21	0.44	0.89	0.011	40	0.56 *
FR Total	-0.18	0.33	0.51	0.70	0.0004	25	0.72 **
DLE	-0.18	-0.17	0.02	0.82	0.906	34	0.54 *
DRE	-0.12	-0.10	0.02	1.23	0.918	76	0.07
Diotic	-0.15	0.23	0.38	0.62	0.002	19	0.71 **
Diotic Adv	0.03	0.19	0.16	0.99	0.381	49	0.25
FR REA	-0.09	-0.07	0.02	1.40	0.945	98	-0.03
Directed REA	0.02	0.03	0.01	1.07	0.976	57	0.36 *
Attention Adv	-0.18	-0.68	-0.50	1.37	0.058	94	0.04

DDdT Condition Correlations

Pearson product-moment correlation analysis were used to analyse whether correlations existed between scores on the various DDdT conditions. Due to the ceiling effects in the adult data, the correlations were conducted on the data from the 62 children. Results are provided in Table 6.

Table 6. Pearson product-moment correlation coefficient (r), analyzed from the arcsine transformed z scores on the various DDdT conditions and difference measures, for the 62 children in the normative data study. Significant correlations are marked with an asterisk (* < 0.05; ** < 0.01; * <0.001).**

Condition	FR LE	FR RE	FR Total	DLE	DRE	Diotic
FR LE	1.00	0.37 **	0.87 ***	0.36 **	0.28 *	0.50 ***
FR RE		1.00	0.77 ***	0.23	0.48 **	0.55 ***
FR Total			1.00	0.36 *	0.43 **	0.62 ***
DLE				1.00	0.32 *	0.32 *
DRE					1.00	0.35 **
Diotic						1.00

The correlation between the diotic condition and the various free recall conditions were highly significant. FR LE ($r = 0.50$, $p = 0.00004$); FR RE ($r = 0.55$, $p < 0.00001$) and FR Total ($r = 0.62$, $p < 0.00001$), as shown in Figure 5.




Figure 5. Scatterplot of the DDdT Free Recall Total z-scores compared to the diotic z scores for the 62 children in the normative data study. The solid line represents least squares regression line.

DISCUSSION

The studies presented here involved the development and evaluation of a new dichotic listening test - the Dichotic Digits difference Test (DDdT). The DDdT was designed with the aim of providing clinicians with additional information as to the validity of a dichotic diagnosis, in clients with suspected CAPD, via the inclusion of a diotic control condition which requires the same response criteria and carries the same memory load as the dichotic free recall task. The inclusion of the diotic condition enabled calculation of a *dichotic advantage* score, measured as the average of the free recall (FR) right and left ear scores (referred to as FR total) minus the diotic score. This difference score provides a means of

1
2
3 assessing genuine dichotic ability by minimizing the impact of cognitive skills on test score.
4
5 The DDdT software utilizes automated scoring and reporting features, and, in a phase one
6
7 study, stimuli were adjusted for equalization of difficulty based on error-rate analysis. This
8
9 equalization makes it possible to pseudo-randomly create multiple lists of close-to-equal
10
11 difficulty.
12

13
14 Normative data was subsequently collected from 62 typically-developing children and 10
15
16 adults. The data was arcsine transformed to achieve a distribution that was closer to a normal
17
18 distribution, and z-scores were calculated. Even when age effects were removed, via
19
20 expression of the results as z-scores, there was a highly significant correlation ($r = 0.62$, $p <$
21
22 0.00001) between performance on the dichotic free recall total and diotic conditions. This
23
24 high correlation presumably reflects the large impact that cognitive skills, unrelated to
25
26 dichotic perception, have on dichotic test results.
27

28
29 In respect to test-retest reliability, the DDdT conditions that improved significantly with
30
31 practice – dichotic free recall LE, RE and total, as well as diotic, are those that would be
32
33 expected to improve if cognitive skills relevant to the test got better. That is, the participant's
34
35 response "strategies" – such as visualizing the sets of numbers - are better the second time
36
37 around. Test scores that we would expect to be less affected by verbal memory skills – DDdT
38
39 directed left ear (LE) and right ear (RE) conditions and dichotic advantage – did not improve
40
41 with practice.
42
43

44
45 Based on the percentage of the variance estimated to be accounted for by measurement
46
47 error, it might at first appear from Table 5 that many of the measurements are of little value,
48
49 with the error percentage varying from close to half up to 98% for 6 of the 10 measures
50
51 shown. One of these measures is the free recall LE condition, which is routinely reported in
52
53 auditory processing literature. Because scores for each ear are based on only 40 scored items,
54
55 scores will have a significant random component that can easily be calculated from the
56
57
58
59
60

1
2
3 binomial distribution. The standard deviation for repeated application of the test, even when
4
5 there are no changes in true ability caused by practice, fatigue, fluctuation in attention, or
6
7 variation in listening strategies is equal to $\sqrt{(P(1-P)/N)}$, where P is the true proportion of
8
9 items correctly repeated and N is the number of scored items. For a test with 40 scored
10
11 digits, and true scores around 80% for example (the average score for left ear free recall for
12
13 an 8 year old), the expected standard deviation is 6.3%. This means that 95% of the time
14
15 scores for a child with this true score will fluctuate over a range from approximately 67% to
16
17 93% (and 5% of the time the fluctuation will be even greater). Of course, practice, fatigue,
18
19 fluctuations in attention and variations in listening strategy all can occur, so an even bigger
20
21 fluctuation than this would be expected in practice. However, these data were obtained from
22
23 typically-developing children. Were children with more aberrant scores to be included, the
24
25 range of raw scores would be greater, and the percent variance accounted by measurement
26
27 error would reduce accordingly.
28
29
30
31

32 It is evident from Figure 4 that the +2 SD limits are wider for the two REA difference
33
34 measures than for either dichotic advantage or attention advantage. This is not surprising as
35
36 the scores from which the REA measures are derived are each based on scores for 40 digits,
37
38 whereas the dichotic and attention advantage measures are derived from scores based on 80
39
40 digits. The REA measures should thus contain greater measurement error. Consistent with
41
42 this, the test-retest correlations are larger for the FR total scores and diotic scores, each of
43
44 which is based on 40 digits. The very large error component of the two right ear advantage
45
46 measures should be a warning against taking measurement results at face value. It is
47
48 common to seize on a larger than average asymmetry in dichotic test results as an indication
49
50 of an underlying problem. Quantifying the right ear advantage with z-scores as we have done
51
52 makes it possible to determine whether an apparent ear asymmetry really does represent a
53
54
55
56
57
58
59
60

1
2
3 problem, or is just the result of the child achieving better than his/her true score in one ear
4
5 and poorer than his/her true score in the other, both due to chance factors.
6
7
8
9

10 11 12 **CONCLUSION**

13
14
15
16 The new dichotic digits difference test presented in this article includes a diotic condition that
17
18 can act as a control to help differentiate children who perform below normal on a dichotic test
19
20 due to a dichotic perception issue from those who perform below normal because of a
21
22 cognitive issue. Interpretation of the control condition results is facilitated by expressing all
23
24 results in age-corrected z-score units, including the differential ability to use dichotic cues
25
26 when they are present. An additional study with both clinical and non-clinical participants
27
28 which investigated the correlations between DDdT conditions, as well as between DDdT
29
30 conditions and various cognitive abilities (memory, attention and intelligence) will be
31
32 described in the article which follows.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Ahmmmed AU, Ahmmmed AA, Bath JR, Ferguson MA, Plack CJ, Moore DR. (2014) Assessment of children with suspected auditory processing disorder: A factor analysis study. *Ear Hear* 35(3): 295-305.
- American Academy of Audiology. (2010) Diagnosis, treatment and management of children and adults with central auditory processing disorder [Clinical Practice Guidelines]. Retrieved from http://audiology-web.s3.amazonaws.com/migrated/CAPD%20Guidelines%208-2010.pdf_539952af956c79.73897613.pdf.
- American Speech-Language-Hearing Association. (2005) (Central) Auditory Processing Disorders [Technical Report]. Available at <http://www.asha.org/policy/TR2005-00043/>.
- Bellis TJ. (2014) The nature of central auditory processing disorder. In: Chermak GD and Musiek FE, Eds. *Handbook of Central Auditory Processing Disorder. Auditory Neuroscience and Diagnosis* (Vol I). San Diego: Plural Publishing, 211-230.
- Bellis TJ. (2003) *Assessment and management of central auditory processing disorders in the educational setting. From Science to practice*. New York: Delmar Learning.
- Cameron S, Glyde H, Seymour J, Dillon H. (2013) Dichotic Digits difference Test (DDdT) (Research Version) [Computer software]. Sydney, NSW: National Acoustic Laboratories.

1
2
3 DeBonis DA. (2015) It is time to rethink central auditory processing disorder protocols for
4
5 school-aged children. *Am J Audiol* 24: 124-136.
6
7

8
9
10 Hiscock M, Kinsbourne M. (2011) Attention and the right-ear advantage: What is the
11
12 connection? *Brain Cogn* 76: 263-275.
13

14
15
16 Kimura D. (2011) From ear to brain. *Brain Cogn* 76: 214-217.
17

18
19
20 Maerlender A. (2010) Short-term memory and auditory processing disorders: Concurrent
21
22 validity and clinic diagnostic markers. *Psychol Schools* 47(10): 975-984.
23
24

25
26
27 Martin J, Jerger J, Mehta J. (2007) Divided-attention and directed-attention listening modes
28
29 in children with dichotic deficits: an event-related potential study. *J Am Acad Audiol* 18(1):
30
31 34-53.
32
33

34
35
36 MATLAB (Release 2013a) [Computer software]. (2013). Natick, Massachusetts: The
37
38 MathWorks Inc.
39
40

41
42
43 Moncrieff DW. (2011) Dichotic listening in children: Age-related changes in direction and
44
45 magnitude of ear advantage. *Brain Cogn* 76: 316-322.
46
47

48
49 Musiek FE, Weihing J. (2011). Perspectives on dichotic listening and the corpus callosum.
50
51 *Brain Cogn* 76: 225-232.
52
53
54
55
56
57
58
59
60

1
2
3 Schmithorst V, Farah R, Keith RW. (2013) Left ear advantage in speech-related dichotic
4
5 listening is not specific to auditory processing disorder in children: A machine-learning fMRI
6
7 and DTI study. *Neuroimage Clin* 3: 8-7.
8
9

10
11 Tomlin D, Dillon H, Kelly AS. (2014) Allowing for asymmetric distributions when
12
13 comparing auditory processing test percentage scores to normative data. *J Am Acad Audiol*
14
15 25:541-548.
16
17

18
19
20 Tomlin D, Dillon H, Sharma M, Rance G. (2015) The impact of auditory processing and
21
22 cognitive abilities in children. *Ear Hear*, doi: 10.1097/AUD.000000000000172.
23
24

25
26
27 Weihing J, Atcherson SR. (2014) Dichotic listening tests. In: Chermak GD and Musiek FE,
28
29 Eds. *Handbook of Central Auditory Processing Disorder. Auditory Neuroscience and*
30
31 *Diagnosis* (Vol I). San Diego: Plural Publishing, 369-404.
32
33

34
35
36 Westerhausen R, Hugdahl K. (2008) The corpus callosum in dichotic listening studies of
37
38 hemispheric asymmetry: A review of clinical and experimental evidence. *Neurosci Biobehav*
39
40 *Rev* 32: 1044-1054.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FIGURE CAPTIONS

Figure 1. DDdT graphical user interface (research version).

Figure 2. Percent of incorrect response for each digit pair combination for the 20 adults in the development study.

Figure 3. Scatterplots of the DDdT regression curves (solid line) transformed back to percentage values (i.e. percent correct): (a) FR LE; (b) FR RE; (c) FR Total; (d) DLE; (e) DRE; (f) Diotic, based on the data from the 62 children and 10 adults in the normative data study. Original data is represented by the blue circles. The dashed line represents the percentage values that correspond to the transformed values that are two standard deviations below the age-dependent mean values.

Figure 4. Scatterplots of performance on the DDdT difference measures (Dichotic Advantage, Free Recall REA, Directed REA and Attention Advantage) by the 62 children in the normative data study. The solid line indicates the mean score as a function of age, the dashed lines show the + and – 2 standard deviation limits.

Figure 5. Scatterplot of the DDdT Free Recall Total z-scores compared to the diotic z scores for the 62 children in the normative data study. The solid line represents least squares regression line.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1
110x135mm (150 x 150 DPI)




Figure 2
165x93mm (150 x 150 DPI)

Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 3a
159x119mm (96 x 96 DPI)

Review




Figure 3b
160x120mm (96 x 96 DPI)

Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 3c
159x119mm (96 x 96 DPI)

Review




Figure 3d
159x119mm (96 x 96 DPI)

Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 3e
159x119mm (96 x 96 DPI)

Review




Figure 3f
159x119mm (96 x 96 DPI)

Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 4a
165x123mm (96 x 96 DPI)

Review




Figure 4b
165x123mm (96 x 96 DPI)

Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 4c
165x123mm (96 x 96 DPI)

Review




Figure 4d
165x123mm (96 x 96 DPI)

Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 5
165x123mm (96 x 96 DPI)

Review

Table 1. Average percent correct scores on the six DDdT conditions as a function of age.

	Age					
	7	8	9	10	11	Adult
FR LE %	79.5	73.2	79.6	88.8	90.6	99.8
FR RE %	88.9	85.0	86.3	91.8	91.7	99.8
FR Total %	84.4	79.1	82.9	90.3	91.1	99.8
DLE %	81.3	86.1	86.5	91.6	94.4	99.8
DRE %	91.6	87.1	91.9	95.0	96.7	99.3
Diotic %	72.1	72.0	77.3	83.2	87.5	96.0

Table 2. Regression coefficients for the exponential formula shown in Equation 2 ($T' = a - b.e^{(-age/c)}$). Also shown is σ , the standard deviation of the residuals (transformed scores minus predicted transformed scores).

Condition	<i>a</i>	<i>b</i>	σ
FR LE	1.572	1.410	0.160
FR RE	1.542	0.972	0.148
FR Total	1.546	1.201	0.120
DLE	1.581	1.180	0.149
DRE	1.565	0.786	0.161
Diotic	1.440	1.201	0.116

1
2
3
4
5
6
7 **Table 3. Regression coefficients for the equation shown in Equation 4. Also shown is σ ,**
8 **the standard deviation of the residuals (difference scores minus predicted difference**
9 **scores).**
10
11
12
13
14
15
16
17

Difference score	<i>f</i>	<i>g</i>	σ
Dichotic advantage	0.247	-0.0155	0.101
REA – free recall	0.390	-0.0317	0.173
REA – directed	0.176	-0.0081	0.182
Attention advantage	-0.003	0.0106	0.121

18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 4. Regression coefficients for the equation shown in Equation 6. Also shown is σ , the standard deviation of the residuals (difference scores minus predicted difference scores).

Difference score	<i>h</i>	<i>i</i>	<i>j</i>	σ
Dichotic advantage	3.10	61.12	3.50	7.60
Right ear advantage – free recall	0	176.20	2.75	9.45
Right ear advantage – directed	0	50.41	3.91	10.48
Attention advantage	4.43	0	-	7.45

Table 5. Average test, retest and retest minus test z-scores and SDs for each DDdT condition and advantage measure for the 30 children who took part in the test-retest reliability study. P-values from repeated measures ANOVAs of test and retest z scores and error variance (%) are also provided. Pearson product-moment correlation coefficient (r) is shown, with significant test-retest correlations marked with an asterisk (* < 0.05; ** < 0.01).

Condition	<i>Test Mean z-score</i>	<i>Retest Mean z-score</i>	<i>Retest-Test Mean z-score</i>	<i>SD</i>	<i>p-value</i>	<i>Error variance (%)</i>	<i>r</i>
FR LE	-0.08	0.30	0.39	1.06	0.055	56	0.36 *
FR RE	-0.24	0.21	0.44	0.89	0.011	40	0.56 *
FR Total	-0.18	0.33	0.51	0.70	0.0004	25	0.72 **
DLE	-0.18	-0.17	0.02	0.82	0.906	34	0.54 *
DRE	-0.12	-0.10	0.02	1.23	0.918	76	0.07
Diotic	-0.15	0.23	0.38	0.62	0.002	19	0.71 **
Diotic Adv	0.03	0.19	0.16	0.99	0.381	49	0.25
FR REA	-0.09	-0.07	0.02	1.40	0.945	98	-0.03
Directed REA	0.02	0.03	0.01	1.07	0.976	57	0.36 *
Attention Adv	-0.18	-0.68	-0.50	1.37	0.058	94	0.04

Table 6. Pearson product-moment correlation coefficient (r), analyzed from the arcsine transformed z scores on the various DDdT conditions and difference measures, for the 62 children in the normative data study. Significant correlations are marked with an asterisk (* < 0.05; ** < 0.01; * < 0.001).**

Condition	FR LE	FR RE	FR Total	DLE	DRE	Diotic
FR LE	1.00	0.37 **	0.87 ***	0.36 **	0.28 *	0.50 ***
FR RE		1.00	0.77 ***	0.23	0.48 **	0.55 ***
FR Total			1.00	0.36 *	0.43 **	0.62 ***
DLE				1.00	0.32 *	0.32 *
DRE					1.00	0.35 **
Diotic						1.00