

CHAPTER 3

LITERATURE REVIEW (B): LEISURE NOISE

Leisure noise

The work presented in this chapter has been published as:

Carter, L., Black, D., Bundy, A., & Williams, W. (2014). The leisure-noise dilemma: hearing loss or hearsay? What does the literature tell us? *Ear and Hearing* 35(5): 491-505.

As first author of the paper "*The leisure-noise dilemma: hearing loss or hearsay? What does the literature tell us?*" I, Lyndal Carter, confirm that I have made the following contributions: devised the study concept, performed the data collection (literature review) with the guidance of the co-authors, and drafted the manuscript in full, in consultation with the co-authors.

Signed.....Date:.....21/7/16....



As supervisor for the candidature upon which this thesis is based, I can confirm that the authorship attribution statements above are correct.

Supervisor Name: Prof Deborah Black

Signed.....Date:.....21/7/16....



Title: The leisure-noise dilemma: hearing loss or hearsay? What does the literature tell us?

Authors: Lyndal Carter¹, Deborah Black³, Anita Bundy³, Warwick Williams^{1,2}

1. National Acoustic Laboratories, Sydney, NSW, Australia.

2. The Faculty of Health Sciences, University of Sydney, New South Wales, Australia.

3. The HEARing CRC, East Melbourne, Victoria, Australia.

Key words: Adolescents, Hearing Loss, Noise-Induced, Leisure, Young adults.

Abbreviations: AEF = auditory evoked magnetic field; ATS = asymptotic threshold shift; CI = cochlear implant; CRC = Cooperative Research Centre; dB = decibels; HI = hearing impaired; HL = hearing level; HTL = hearing threshold level; ISO = International Organisation for Standardisation; L_{Aeq} = long-term equivalent average sound level; $L_{Aeq,8h}$ = exposure level, 8 hour equivalent continuous A-weighted SPL; MEG = magnetoencephalography; NAL = National Acoustic Laboratories; NH = normal (non-impaired) hearing; NHANES = National Health and Nutrition Examination Survey; NHES = National Health Examination Survey; NIHL = noise-induced hearing loss; NIPTS = noise-induced permanent threshold shift; NITS = noise-induced threshold shift; NHMRC = National Health and Medical Research Council; NSW = New South Wales; OAE = otoacoustic emission; OHS = Office of Hearing Services; Pa^2h = Pascal squared hours; PCP = personal cassette player; PHP = personal hearing protector; PSP = personal stereo player; PTA = pure tone audiometry; PTS = permanent threshold shift; QoL = quality-of-life; SPL = sound pressure level; TTS = temporary threshold shift; SCENIHR = Scientific Committee on Emerging and Newly Identified Health Risks; WHO = World Health Organization. 4FAHL = four frequency average hearing level of 500, 1000, 2000, & 4000 Hz.

Definitions:

1. Loud sound encountered during everyday leisure activities is variously referred to as “leisure noise”, “social noise” (Smith et al., 2000), and “sociacusis” (Ward, 1976; Medical Research Council (MRC), 1986; Yaremchuk et al. 1997). The term “leisure noise” is used in this review.

2. The terminology associated with “hearing loss” also varies in the literature and among authorities (e.g., standards organisations and professional bodies). Whatever the nomenclature used, the following concepts should be differentiated: (1) *Threshold shift* (or, *threshold impairment*), that is, deviation, or worsening of individual hearing threshold levels (HTLs) from a baseline or, alternately; the HTL of an individual (or group) in relation to an accepted audiometric standard (WHO, 1980); 2) *Noise-induced threshold shift (NITS)*, that is, threshold shift attributable to noise exposure alone and; (3) *Hearing handicap* (or, *hearing disability/hearing impairment*), that is, the individual disadvantage in everyday life imposed by threshold shift, particularly in terms of understanding conversational speech (ISO, 1990 p 3). “Hearing loss” generally refers to threshold shift in this review.

Parts of this article were presented as an oral poster at the Audiology Australia XX National Conference. 1 – 4 July 2012. Adelaide, Australia.

Acknowledgements: The authors acknowledge the extensive contribution of Joy Fischer (National Acoustic Laboratories Librarian) in sourcing and obtaining the material for this review, and Elizabeth Beach, Harvey Dillon, John Macrae and Denise Macoun for their comments on the draft manuscript.

ABSTRACT

The authors undertook a review of the literature, focusing on publications describing the following:

- 1) Pure tone threshold data for adolescents/young adults;
- 2) Measurements/estimates of noise exposure from leisure activities; and
- 3) The relationship between HTLs and leisure-noise exposure.

There is a large volume of published materials relevant to these topics, and opinion among authors regarding the relationship between leisure-noise exposure and HTLs varies significantly. At one extreme is the view that the effects of leisure noise are minimal. The opposing belief is that as a direct result of leisure-noise exposure, significant HTL shifts and possibly significant hearing disability are occurring in a large (and increasing) proportion of young people. It has been claimed that behaviours relating to leisure noise are “as threatening to young people’s health as more traditional risk behaviours” (Bohlin & Erlandsson, 2007, p. 55). This view has been reiterated by the popular media. This review revealed that while sufficient data confirm that some leisure pursuits provide potentially hazardous noise levels, the nature of the exposure-injury relationship for leisure noise is yet to be determined. Specific information about the quality-of-life impacts of threshold shift related to leisure-noise exposure is also lacking. The scope and limitations of a large sample of relevant publications and an overview of the methodological issues in this area of research are briefly presented. Considerations for future research are raised.

INTRODUCTION

There is no doubt that exposure to noise (i.e., loud sound) of enough intensity over sufficient periods of time can result in temporary and permanent HTL shifts (Miller, 1974; Mills, 1975; Ylikoski et al., 2001; Zhao et al., 2010). Underlying pathological processes arising from noise exposure have been described in much detail, (e.g., Nicotera et al., 2001; Wang et al., 2002; Kujawa & Liberman, 2009; Makary et al. 2011; Op de Beeck et al, 2011). There appear to be a number of possible biological mechanisms (Henderson et al., 1993) and large inter-individual variations in susceptibility to noise injury (Quaranta et al., 2001) that are still not fully understood. Auditory system damage may precede observable changes in the pure tone audiogram (Axelsson, 1991; Axelsson et al., 1994; Smith et al., 2000; Kujawa & Liberman,

2009; Jin et al., 2013). Carter et al. (1978; citing a study by Bienvenue et al., 1976) noted that noise has been shown to have a temporary effect on the loudness difference limen. West and Evans (1990) investigated frequency resolution abilities, reporting that participant groups “more exposed” to amplified music had wider bandwidths than less exposed groups under some conditions. Okamoto et al. (2011) reported a study of magnetoencephalographic (MEG) responses of long-term users and non-users of personal stereo players (PSPs). Both groups performed equally on standard audiometric evaluations (including pure tone audiometry [PTA]); however, significantly broadened population-level frequency tuning in a group of long-term users under a specific listening condition was observed using MEG. Kumar et al. (2012) observed deterioration in temporal processing and speech processing abilities of individuals exposed to occupational noise, with HTLs better than 25 dB hearing level (HL) in the octave frequencies between 250 and 8000 Hz. Studies have also revealed an association between tinnitus and leisure-noise exposure, even in the presence of clinically unremarkable HTLs (Davis et al., 1998; Tin & Lim, 2000; Holgers & Petterson, 2005; Beach et al., 2013a).

In 1975, the International Organisation for Standardization (ISO) published the first edition of its standard describing the statistical relationship between occupational noise exposure and noise-induced permanent threshold shift (NIPTS) in people of various ages – ISO 1999 (ISO, 1990). This description is based on data from a number of earlier, cross-sectional studies of workers’ pure tone hearing thresholds. ISO 1999 provides the first reliable description of “noise-exposure – hearing effect” (Williams, 2011, p. 13) or the dose-response relationship between *occupational* noise exposure and pure tone threshold shift.

The association between noise exposure and noise injury is regarded as stronger for occupational noise than leisure noise (Hidecker, 2008). It is believed, however, that

technological advances, particularly the proliferation of PSPs, have led to dramatically increased leisure-noise exposure (Zhao et al., 2010; Levey et al. 2011), with a concomitant increase in risk for young people. In the 1960s, “the damaging effects of rock and roll music on hearing” (Rintelmann & Borus, 1968, p. 57) were a new cause for concern. A large body of literature concerned with the possible relationship between leisure-noise exposure and hearing threshold shift has since amassed – however there is still a lack of consensus about the extent of the risk. At one extreme “...there is a concern we may be facing an epidemic of hearing impairment” (Agrawal, 2008, p. 1522). The opposing viewpoint is that the effects of leisure noise are “slight” (Carter et al, 1984).

The leisure-noise issue has received significant media attention (Carter et al., 1978; Smith et al., 2000; Schlauch & Carney, 2011), with the popular press tending towards alarmist headlines, which Hètu and Fortin (1995) suggested denote a disapproving attitude towards particular leisure activities (e.g., rock music). A critical attitude is also discernible in some scientific publications. Maassen et al. (2001, p. 4), for example, commented that “*A “techno freak” subjecting himself to loud music via a PCP [personal cassette player] endangers his ears in the same way as a worker in a steel factory using no ear protection.*” Hètu and Fortin (1995) suggested that such assertions have received a wide and largely accepting audience. Further, leisure-noise exposure differs from occupational exposure with respect to the fact that individuals participate voluntarily in noisy recreational activities according to their own preferences. The restriction of preferred activities may be considered a legitimate “cost” (Hill, 1965; Phillips and Goodman, 2004), a factor which appears to have received relatively little attention in either public or scientific commentary on this issue.

In recent years, the National Acoustic Laboratories (NAL) has conducted a range of studies aiming to quantify leisure-noise sources and patterns of exposure, estimate community risk of noise injury from everyday non-work activities, and determine the prevalence of hearing threshold impairment in the younger Australian population. This review was undertaken with no previous intent to justify a particular position in the leisure-noise debate but rather with the aim of providing an objective frame of reference for disseminating recent NAL findings and for considering methodological “best practice” for ongoing research.

MATERIALS AND METHODS

An extensive literature review was conducted using commonly accessed Internet search methods (particularly, PubMed, Google Scholar and MEDLINE) and scrutiny of the reference lists of peer-reviewed publications considered to be of high relevance. Search terms included: *hearing threshold levels, leisure noise, music and hearing, recreational noise, prevalence of hearing loss, and personal stereo players*. Title selection and review were performed only by the first author. Only titles in English or with a translated abstract were considered. No meta-analysis was performed.

About 737 titles of interest (including peer-reviewed publications, referenced conference abstracts, and postgraduate theses) were identified in this review. It was evident that the number of publications relevant to this topic has increased steadily over the past two decades, as noted by other authors (Morata, 2007; Zocoli et al., 2009). Figure 3.1 shows the number of titles identified by decade (relating to leisure noise and hearing), from the 1940s until the end of the first decade of this century.




Figure 3.1: Publications by decade (leisure noise and hearing).

It was beyond the scope of this review to fully appraise all of this material. Because several substantial reviews have been previously published (e.g., Mills, 1975; MRC, 1986; Clark, 1991), more emphasis was placed on recent material. The total number of publications for review was also reduced as follows: articles that primarily addressed attitudes toward noise and descriptions of hearing loss prevention interventions or those providing hearing threshold data for very young children (particularly those obtained in screening programs) were eliminated. Articles pertaining primarily to the effects of noise exposure on otoacoustic emissions (OAEs) and articles relating specifically to occupational noise-exposure (apart from musicians) were also excluded. Articles pertaining to professional classical musicians were excluded. Some articles containing data about employees in amplified music venues, however, were included, in view of the overlap between occupational and recreational exposure for rock and pop music. Articles relating to firearms use were excluded, on the basis that the serious threat of noise injury from firearms use (Clark, 1991) is not controversial. A total of 265 articles were reviewed in full. The content of these key articles, and supplementary details from another 145 topical abstracts (e.g., articles that were of interest but were not in the English language, or could not be obtained), provided the basis for the following commentary.

RESULTS

Sound pressure level (SPL) measurement of leisure-noise sources

Over 100 reports of this type were identified in this review. Results obtained using one or both of the following methods have typically been described:

1. Sampling the sound pressure level (SPL) using a sound level meter at a fixed position(s) – either in situ (at the activity site/venue), or in simulated laboratory conditions.
2. Measuring the SPL in real-life situations over a period of time, using individually worn noise exposure meters (dosimeters) – a technique which has become more feasible and more informative in recent years.

The difficulty in directly comparing the findings of different studies, which have employed a variety of specific methods and metrics, is noteworthy. Weaknesses in the reported data for non-occupational noise, such as inconsistencies in noise level documentation, and the tendency to focus on the highest possible exposures during the noisiest activities, have also been noted (Neitzel et al. 2004).

Early reports (dating from the 1960s) were reviewed by Mills (1975). Mills' article presents the SPLs of firecrackers, model aeroplanes, snow-mobiles and firearms (including toy guns). A later review by Clark (1991) presented maximum SPLs of a variety of common recreational, domestic (e.g., food blender, vacuum cleaner etc.), hobby, transportation, and firearm noise sources. In addition, Clark (1991) provided an overview of studies (conducted in the 1970s and 80s) of rock concerts and the emerging "discotheque". Various reports of SPLs at rock music performances were subsequently published (e.g., Drake-Lee, 1992; Yassi et al., 1993).

Clark (1991) also presented SPL data for early models of PSP, sourced from eight different publications (1972 – 1985). There have been at least three subsequent reviews of published PSP research which provided output level data (Smith et al., 2000; SCENIHR, 2008; Punch et al., 2011). Most recently, Portnuff et al. (2013) reported on PSP outputs, concluding that a “small but significant” percentage of PSP users reported exposure sufficient to increase the risk of noise injury. This conclusion is consistent with those of earlier reviews, although, as noted by Portnuff et al., much higher estimates of risk have been given by some authors. For example, Levey et al. (2011) estimated that 58.2% of participants in their study ($N = 189$) exceeded recommended workplace exposure levels. Measurements in the Levey et al. study were, however, made in a single, high-background noise environment.

The level of noise emitted by children’s toys has also been a subject of recurring interest. Subsequent to the reports described by Mills (1975), Yaremchuk et al. (1997) measured the level of 25 toys (e.g., bicycle horns, toy guns, toy tools, telephones and musical instruments). More recently, Bittel et al. (2008) reported the output levels of 24 commercially-available toys, noting that many toys exceeded recommended safety standards. Mahboubi et al. (2013) reported an experiment in which more than 200 toys were screened for loudness and 90 analysed under controlled conditions. They concluded that acoustic trauma from children’s toys continues to be a potential risk.

The level of noise generated by crowds at public events such as sports matches and rock concerts has received little attention but is also relevant. Opperman et al. (2006) described measures of stadium noise published in 1987 and observations of the contribution of crowd yelling and screaming to overall noise levels in their own study. Beach et al. (2013b) also make reference to crowd situations, such as sporting events, in their discussion.

In the last decade, SPL measures have been reported for a range of other leisure activities, such as electronic arcade games (Mirbod et al., 1992), car stereos (Ramsey & Simmons, 1993), air shows (Pääkkönen et al., 2003), Korean karaoke singing (Park, 2003), aerobics classes (Torre III & Howell, 2008; Beach et al., 2013c), auto racing (Rose et al., 2008; Kardous & Morata, 2010), indoor hockey (Cranston et al., 2013), and marching bands (Jin et al., 2013). Results of some of these studies indicate that when typical activity durations are taken into account, the risk to patrons may not be significant (e.g., Ramsey & Simmons, 1993; Pääkkönen et al., 2003; Rose et al., 2008). Other studies indicate that typical participation in some environments may place individuals at risk, for example, aerobics classes, stock car racing tracks, indoor hockey arenas and karaoke singing venues.

“Daily life” measures, obtained using individual dosimetry, have been reported by several authors (e.g., Neitzel et al. 2004; Flamme et al., 2012; Beach et al., 2013b). In these studies, amplified music stands out as a concerning source of leisure-noise exposure. Beach et al. (2013b) described a contemporary inventory, referred to as the “NOISE” (non-occupational incidents, situations and events) database, in which over 500 dosimetry samples obtained since 2008 have been indexed and categorised for general reference. A number of the loudest samples are contained under the categories “attendance at entertainment venues” (which include karaoke events, nightclubs, dance clubs and discos) and “arts and cultural activities” (which include live music performances, popular music concerts and music festivals).

Overall, there is reasonable agreement among authors that some leisure activities (particularly shooting and amplified music listening) provide SPLs which would be of safety concern in industrial settings (Tambs et al., 2003; Zhao et al., 2010). Although the levels of some leisure

activities are analogous to those encountered in occupational settings, it is important to reiterate that the damaging effects of noise depend not only on intensity but on the duration and pattern of exposure and possibly on other individual susceptibility factors. To date, risk estimates for leisure-noise exposure have been based on dose-response relationships observed in the industrial setting (e.g., ISO 1999, 1990; ISO 1999, 2013), which assume continuous 8 hour daily exposure over many years (Hètu & Fortin, 1995). Strasser et al. (2003) cautioned that rating sound exposures by energy equivalence alone can lead to very misleading assessments of their actual physiological costs.

Apart from the fact that noise exposure during leisure is typically less frequent and intense than that encountered in the workforce, there is also the possibility that non-industrial sources (because of their unique physical characteristics) may have distinctive effects on the auditory system. Most music, for example, has a greater variation in spectral content and intensity and a greater spread of energy over time, compared with typical industrial noise sources (Turunen-Rise et al., 1991). It has been suggested that the intermittent nature of music may reduce the risk of noise-induced hearing loss (NIHL) occurring (Jin et al., 2013), and that exposure to some types of moderate-level noise may have a “conditioning” or protective effect on the cochlea (McFadden et al., 1997; Niu, 2002). The “heavy metal” genre, however, has been reported as more similar in effect to industrial sources (Strasser et al., 1999).

Studies exploring the relationship between leisure-noise exposure and HTL

Obviously, it is ethically impossible to determine the noise-injury relationship in human participants via direct experimental means. The following section describes the six main methodological approaches that have been applied to this research question. The main studies identified in each of these categories are listed in Tables 3.1 - 3.6.

Pre-exposure/Post-exposure assessments

The effects of leisure-noise exposure have been investigated experimentally using pre-exposure and post-exposure audiometry (and/or other measures, such as OAEs), to look for evidence of post-exposure shift, then recovery of HTLs (i.e., temporary threshold shift, or TTS). Table 3.1 lists the methods and findings of 19 investigations of this type, conducted from the late 1960s to date.

Table 3.1: Studies of TTS.

Authors	Participants	Method/noise source	Conclusion
<i>Rintelmann and Borus (1968)</i>	<i>N</i> = 52 United States (U.S.) 18–20 yrs	PTA (pre/post live rock music)	Concern seems unwarranted
<i>Reddell and Lebo (1972)</i>	<i>N</i> = 43 U.S. rock musicians Mean age 22 yrs	PTA (pre/post hard rock music)	TTS observed in musicians and some listeners
<i>Axelsson and Lindgren (1978)</i>	<i>N</i> = 83 Swedish pop musicians and listeners	PTA (pre/post pop music)	Less TTS in musicians than listeners
<i>Lindgren and Axelsson (1983)</i>	<i>N</i> = 10 Swedish teenagers	PTA (pre/post 10 min laboratory stimuli)	Noted differences in TTS with musical vs. nonmusical stimuli
<i>Lee et al. (1985)</i>	<i>N</i> = 16 U.S.	PTA (pre/post 3 hr PCP exposure)	6/16 showed TTS All recovered after 24 hours
<i>Clarke and Bohne (1986)</i>	<i>N</i> = 6 U.S. rock concert attendees	PTA (pre/post rock concert)	5/6 showed TTS
<i>Swanson et al. (1987)</i>	<i>N</i> = 20 U.S. undergraduate students	PTA, tympanometry, acoustic reflex thresholds (pre/post laboratory noise & music)	Relationship found between TTS and music preference (greater for disliked music)
<i>Hellström et al. (1988)</i>	<i>N</i> = 21 Swedish PCP/speaker listeners	Bekesy (pre/post 1 hour of PCP use)	Most had only “discrete” TTS, despite levels of 91–97 dB
<i>Drake-Lee (1992)</i>	<i>N</i> = 5 United Kingdom (U.K.) heavy metal players 25–37 yrs	PTA (pre/post rock concert)	TTS noted in all but one musician (who used PHP)
<i>Yassi et al. (1993)</i>	<i>N</i> = 22 Canadian 18–40 yrs	PTA (pre/post rock concert)	81% showed TTS of 10 dB or more
<i>Vittitow et al. (1994)</i>	<i>N</i> = 12 U.S.	PTA (pre/post music and cycling)	Greater TTS for noise and exercise condition than noise alone

<i>McCombe et al. (1995)</i>	<i>N</i> = 18 U.K. motorcyclists	PTA (pre/post 1 hour motorcycle ride)	Significant TTS found at 0.25–2 kHz
<i>Strasser et al. (1999)</i>	<i>N</i> = 10 German 18–30 yrs	PTA (pre/post laboratory music vs. industrial and white noise)	Demonstrated TTS with all sources Least effect with classical music Industrial noise and heavy metal music showed similar effect
<i>Mazelova et al. (2001)</i>	<i>N</i> = 12 Czech 18–25 yrs	PTA, Bekesy high resolution, OAE (pre/post laboratory amplified music)	Demonstrated changes in all measures except gap detection
<i>Nassar (2001)</i>	<i>N</i> = 28 U.K. Mean age 21 yrs	PTA (pre/post aerobics class)	Exposed group showed TTS, control group slightly improved HTLs (practice effect?)
<i>Sadhra et al. (2002)</i>	<i>N</i> = 14 U.K. university student bar employees 20–40 yrs	PTA (pre/post bar/discotheque music)	13/14 showed TTS
<i>Emmerich et al. (2002)</i>	<i>N</i> = 34 German 18–24 yrs	PTA & AEF (pre/post discotheque music)	TTS found in all subjects and AEF latency shifts
<i>Opperman et al. (2006)</i>	<i>N</i> = 29 U.S. 17–59 yrs	PTA (pre/post concert – amplified music)	64% of unprotected listeners showed TTS, 27% of those using earplugs
<i>Keppler et al. (2010)</i>	<i>N</i> = 21 Belgian 19–28 yrs	PTA & OAE (pre/post high-level MP3 pop/rock music)	Changes in PTA and TEOAE in exposed group No significant changes in DPOAE
<i>Tam et al. (2013)</i>	<i>N</i> = 12 Australian 19–28 yrs	PTA & OAE (pre/post MP3 music)	Significant increase in 6 kHz HTL and significant reduction in some DPOAE and TEOAE amplitudes post exposure

AEF, auditory evoked magnetic field; DP, distortion product; OAE, otoacoustic emission; PCP, personal cassette player; PHP, personal hearing protector; PTA, pure-tone audiometry; TE, transient evoked; TTS, temporary threshold shift.

Many of the studies listed in Table 3.1 reported positive findings. However, the relationship between TTS and permanent threshold shift (PTS) is still debated (Quaranta et al., 2001; Zhao et al., 2010). Consequently, although studies of this type are of interest, they do not provide conclusive information about the lasting effects of leisure noise on hearing thresholds. It is also possible that even when HTLs recover, lasting physiological changes have nevertheless occurred.

Retrospective cohort studies

In a number of investigations HTLs, and/or other indicators (e.g., OAEs), have been examined in groups of individuals voluntarily exposed to specific leisure-noise sources versus similar, non-exposed (control), participants.

Table 3.2: Retrospective cohort studies.

Authors	Participants	Exposure source	Findings
<i>Hansen and Fearn (1975)</i>	<i>N</i> = 505 U.K. students	Pop music	PTA: Small but statistically significant difference between case & controls
<i>Fearn (1981)</i>	<i>N</i> = 367 U.K. school children	Amplified pop music	PTA: Statistical analysis not presented (differences in order of a few dB)
<i>West and Evans (1990)</i>	<i>N</i> = 60 U.K. 15–23 yrs	Amplified music	Bekesy audiometry and frequency resolution: “Trend” towards wider bandwidths in the exposed
<i>Jorge Junior (1993) cited by Zocoli et al. (2009)</i>	<i>N</i> = 958 Brazilian teenagers	PSPs	PTA: No significant differences
<i>Schmidt et al. (1994)</i>	<i>N</i> = 133 Dutch Music students & controls	Classical music	PTA: No significant difference

<i>Meyer-Bisch (1996)</i>	<i>N</i> = 1364 French ~15–25 yrs	Discotheques, PSPs, rock concerts	PTA: No significant differences for discotheque exposure Small (~2–4 dB) but significantly significant differences comparing controls and the “intensively” exposed for PSP and rock concert exposure
<i>Mostafapour et al. (1998)</i>	<i>N</i> = 50 U.S. college students	PSPs	PTA: No significant differences
<i>Peng et al. (2007)</i>	<i>N</i> = 120 Chinese University students	PSPs	PTA: Statistically significant differences (~3–5 dB) reported

MRC (1986) noted that earlier studies of this type revealed differences in hearing thresholds of no more than a few decibels in noise-exposed versus non-exposed groups. Findings of studies published in the following decade are similarly undramatic (e.g., Schmidt et al., 1994; Meyer-Bisch et al., 1996; Mostafapour et al., 1998). One of the largest studies (*N* = 958) to date (Jorge Junior, 1993), revealed no significant differences between HTLs of PSP users and non-users. More recently, Peng et al. (2007) reported significant differences in HTLs for conventional and extended range audiometry (10 – 20 kHz) between PSP users and non-user controls (*N* = 120/30 respectively). Peng et al. (2007) concluded that 34 out of 240 ears tested (14.1%) showed evidence of hearing loss.

Cross-sectional studies of HTL

A number of cross-sectional audiometric studies similar in size to the cohort studies described above have been undertaken. A summary of seven studies with $N < 500$ is listed in Table 3.3.

Table 3.3: Cross-sectional studies (n < 500) which include comments on leisure-noise exposure.

Authors	Participants	Assessment method	Effect of leisure noise?
<i>Carter et al. (1978)</i>	$N = 231$ Australian university students	PTA; survey – occupational and recreational exposure	No
<i>Lees et al. (1985)</i>	$N = 60$ Canadian 16–25 yrs	PTA; survey – occupational and recreational exposure	Yes. 40% prevalence rate of hearing loss (but insufficient to cause hearing disability)
<i>Ahmed et al. (2007)</i>	$N = 24$ Canadian university students	PTA (PSP users only); survey	No evidence of early hearing loss
<i>Kim et al. (2009)</i>	$N = 490$ Korean adolescents 13–18 yrs	PTA Interview (PSP use)	No relationship between HTL and daily use, however 4000 Hz ‘elevated’ in 24 participants with highest exposure
<i>Martinez-Wbaldo et al. (2009)</i>	$N = 214$ Mexican teenagers	PTA; Survey – noisy activities at school and leisure	“Moderate” association between leisure noise and hearing loss. 20% prevalence rate of loss
<i>Zocoli et al. (2009)</i>	$N = 245$ Brazilian 14–18 yrs	PTA; Survey – noisy leisure activities	No
<i>Le Prell et al. (2011)</i>	$N = 56$ U.S. college students	PTA; Survey – risk factors	“Statistically reliable relationship” between HTL and PSP use in males only

Among these studies, there are an equal number of positive and negative findings. Relatively small sample sizes, and use of convenience samples in smaller cross-sectional and experimental studies, may affect generalisability of results. In addition, it seems possible that experimental work of this type could be subject to publication bias, that is, studies with positive results are more likely to be accepted for publication than those with null or negative results.

Results of a number of larger cross-sectional studies ($N = \sim 500 - 2000$) have also been reported. Ten examples identified in this review are listed in Table 3.4. Only two of these studies suggest an association between HTLs and leisure-noise exposure (Costa et al., 1988; Cone et al., 2010). Costa et al. (1988) reported a higher incidence of high-frequency hearing loss in males than females, and speculated that on the grounds that males typically engage in noisier activities than females, these high-frequency losses may thus be noise-induced. Costa et al. (1988), however, noted that methodological issues, such as cerumen occlusion or collapsing canals, may have affected findings. Apparently, information regarding participant noise exposure was not obtained in this investigation.

Table 3.4: Cross-sectional studies (n ≥ 500) including comments on leisure-noise exposure.

Authors	Participants	Assessment method	Effect of leisure-noise?
<i>Strauss et al. (1977)</i>	N = 1300 German	PTA? [German article]	No
<i>Axelsson et al. (1981)</i>	N = 538 Swedish 17–20 yrs	PTA	No
<i>Carter et al. (1982)</i>	N = 944 Australian 16–20 yrs	PTA; ENT exam	No
<i>Buffe et al. (1986)</i> Article in French cited by <i>Petrescu (2008)</i>	N = 51,726 French 18–25 yrs	PTA; medical exam; noise history	No real correlation between music exposure and HTL (Noted professional DJs had higher HTLs)
<i>Costa et al. (1988)</i>	N = 2264 Swedish 7, 10, 13 yrs	PTA screen (no exposure data or tympanometry)	Yes, on basis that HF loss more common, and males more affected than females
<i>Axelsson (1994)</i>	N = 500 Swedish 18 yrs	PTA	No
<i>Haapaniemi (1995)</i>	N = 687 Finnish 6–15 yrs	PTA; ENT exam; survey	No
<i>Cone et al. (2010)</i>	N = 6591 Australian school children year 1 & 5	PTA screening	Reported PSP use as a risk factor, but most significant factor = NICU admission
<i>Twardella et al. (2011)</i> [German]	N = 2240 German Students grade 9	PTA; medical exam; questionnaire	Non-occupational risk factors identified: firearms, chain saws and power tools
<i>Carter (2011)</i>	N = ~ 1420 Australian 11–35 yrs	PTA; interview; questionnaire	No (preliminary analysis)

DJ, disc jockey; ENT, ear, nose and throat specialist; HF, high frequency; NICU, neonatal intensive care unit; PSP, personal stereo player; PTA, pure-tone audiometry.

Cone et al. (2010) tested the hearing of a large group of elementary school children (N = 6591).

Sensorineural hearing loss was identified in 0.88% (55 of a total of 6581 children assessed).

The use of PSP players was reported to be significantly higher in the group identified with

hearing loss compared with the group with normal hearing. The greatly disparate number of affected versus non-affected cases, however, may cast doubt on Cone et al.’s conclusion that PSP use may be a risk factor for hearing loss. Further, Cone et al. (2010) noted that of the children reported to use PSPs, there was no difference in the reported hours of use per day, or parents’ reports of playing the “stereo too loud” in the normally hearing versus sensorineural hearing loss groups. Overall, the available evidence from the larger cross-sectional studies identified in this review does not suggest a compelling association between leisure-noise exposure and HTLs.

Longitudinal studies

The present review identified only four studies providing serial audiometric data (with various retest intervals) and leisure-noise exposure data for young (mainly teenage) subjects. The details of these studies are listed in Table 3.5.

Table 3.5: Longitudinal studies.

Authors	Participants	Assessment method	Effect of leisure-noise?
<i>Roche et al. (1977, 1979, 1982)</i>	<i>N</i> = 1100 U.S. Main sample 6–18 yrs	PTA over 5 yrs; survey; dosimetry	No significant associations between HTL and noise exposure scores
<i>Carter et al. (1984)</i>	<i>N</i> = 141 Australian 10–12 yrs (at 1 st assessment)	PTA x 2 (retest 6–8 years after baseline)	No significant HTL shifts
<i>Biassoni, et al. (2005); Serra et al. (2005)</i>	<i>N</i> = 173 Argentine 14–17 yrs	PTA annually for four years	Significant downward shift in HTLs for frequencies above 8000 Hz
<i>Jin et al. (2013)</i>	<i>N</i> = 698 U.S. university band members ≤ 25 yrs	PTA; OAE; survey (re-test 3–4 months after baseline)	No significant bilateral HTL shifts reported

A series of reports by Roche et al. (1977, 1979, 1982) described a five-year longitudinal hearing survey (Fels study). Roche et al. (1979) reported no statistically significant associations between noise scores (derived from noise exposure histories) and HTL. For group mean threshold data, however, significant differences were observed for groups with particular exposures (specifically, power tools, Hi Fi, loud TV, and exposure to farm machinery) relative to groups reporting no exposure. Roche et al. (1979) concluded that a longer surveillance period was required for a more effective analysis to be made. Carter et al. (1984) reported no significant shift in HTLs over a 6- to 8-year test period. Biassoni et al. (2005) and Serra et al. (2005) reported a significant downward shift in HTLs, confined to test frequencies above 8000 Hz. However, the lack of normative data for the extended high-frequency range makes this finding difficult to interpret in isolation. Further, Schmidt et al. (1994) suggested that inter-subject variations are greater in extended range audiometry compared with conventional audiometry, and that ageing effects may be present for the very high frequencies even in relatively young people. Schmidt et al. concluded that high-frequency audiometry cannot serve as an early indicator of the traumatic effects of noise. Jin et al. (2013) studied the hearing of a group of U.S. university marching band members and age-matched controls. The period between baseline test (pre-band camp) and follow up (re-test) was 3 to 4 months, with subsequent assessment annually (during band camp). No significant bilateral hearing threshold shifts were reported.

Population surveys of HTL

Several retrospective analyses of HTL data from large population surveys have been published (Niskar et al., 1998; Niskar et al., 2001; Hoffman et al., 2010; Shargorodsky et al., 2010; Henderson et al., 2011). The HTL data analysed were obtained in the U.S. National Health and Nutrition Examination Surveys (NHANESs) which provide ongoing, broad health

surveillance (Flamme et al., 2012). NHANES data sets and experimental protocols are publicly available (Centers for Disease Control and Prevention, 2013) and contain HTL values for large numbers of young participants (e.g., $N = \sim 5000$ for 6- to 19-year-old cohorts of NHANES III).

Niskar et al. (1998, 2001) were the first to publish analyses of NHANES III data, (1988 – 1994) with the aim of estimating the prevalence of hearing loss in the young U.S. population. Niskar et al. (1998) reported that “14.9% of U.S. children have low-frequency or high-frequency hearing loss of at least 16 dB hearing level in one or both ears” (p. 1071). Subsequently, the same authors (Niskar et al., 2001) published an alternate analysis of the same data, concluding that “12.5% of US children aged 6 to 19 years (approximately 5.2 million) are estimated to have noise-induced threshold shift (NITS) in one or both ears”, and stated that “*These findings suggest that children are being exposed to excessive amounts of hazardous levels of noise, and children’s hearing is vulnerable to these exposures*” (p. 40). Subsequently, Shargorodsky et al. (2010) published an analysis of the same NHANES III data, and later NHANES data (2005 – 2006), reporting a prevalence rate of hearing loss of 19.5% among 6- to 19-years-olds for the 2005 to 2006 cohort. As Schlauch and Carney (2011) acknowledge, as the first of their kind, the publications of Niskar et al. are important. However, there are important limitations which cast doubt of the appropriateness of these prevalence estimates. These factors are explained in detail later in this review (see Discussion section).

Hearing survey data from larger population health studies have also been used to look for changes in hearing loss prevalence in populations over time, testing the assumption that technological and social changes have resulted in increased leisure-noise exposure. The conclusions reported are inconsistent. For example, Hoffman et al. (2010) compared National Health Examination Survey I 1959 to 1962 with NHANES 1999 to 2004 data, reaching the conclusion that “*Americans hear as well or better today compared with 40 years ago*”

(Hoffman et al., 2010 p. 725). Shargorodsky et al. (2010) compared NHANES III 1988 to 1994 and NHANES 2005 to 2006 data, concluding that the prevalence of hearing loss among U.S. adolescents aged 12 to 19 years *increased* from 14.9% in 1988 to 1994 to 19.5% in 2005 to 2006. The authors note, however, that the “majority of the hearing loss was slight” (Shargorodsky et al., 2010, p. 775) and most cases were unilateral. Schlauch and Carney (2012) suggested that methodological differences between the two study periods (e.g., different tester qualifications) could affect the estimated hearing loss prevalence. Using slightly different exclusion criteria and definitions of pure-tone average, Henderson et al. (2011) compared the same NHANES data sets as Shargorodsky et al. (2010), reporting no increase in prevalence.

Several reports have also been published comparing the hearing thresholds of other cohorts, also with mixed conclusions. Persson et al.’s (1993) analysis of audiograms of 18- to 19-year-old Swedish military conscripts (obtained between 1969 and 1977) suggested improvement in hearing thresholds across time, which, it was speculated, may reflect improvements in general otological management after the 1950s to 1960s. Axelsson et al. (1994) referred to a 1988 report on Swedish military conscripts (Borchgrevink, 1988) which concluded that the incidence of hearing loss had “doubled” since 1981. Rabinowitz (2006b) reported that U.S. Army data from a similar period (1974 – 1989) showed a decrease in prevalence of hearing loss of army recruits. To date, studies of prevalence trends are few, and whether the rate of NIHL is on the rise seemingly “remains controversial” (Rabinowitz et al., 2006b, p. 369). Further, Lutman and Davis (1994) emphasised that, given all the possible factors involved, it would generally be unsurprising to find substantial variation between different sets of audiometric results, even when comparing large studies (where only small differences would be otherwise expected from statistical uncertainty).

Case reports

The current review identified only two articles containing case reports. First, McMillan and Kileny (1994) presented a single case study of hearing loss documented in a child exposed to an impulse noise from a bicycle horn. The second article, Brookhouser et al., (1992) includes five case studies of young people diagnosed with NIHL (NIHL was assumed on the basis of exposure history provided by the child or others, and the absence of other plausible aetiologies).

DISCUSSION

This review confirmed that PTA has remained the “test of choice” in leisure-noise research. However, Schlauch and Carney (2011, 2012) and others (Green, 2002) have identified notable limitations in: (1) the use of audiometric surveys in general, and (2) the particular analytical techniques employed by some authors in this field of inquiry (e.g., Niskar et al., 1998, 2001; Shargorodsky et al., 2010). Nevertheless, it was evident that the publications of Niskar et al. (1998, 2001) and Shargorodsky et al., (2010) are much cited in support of the position that NIHL in young people is a significant problem (e.g., Chung et al., 2005; Tharpe & Sladen 2008; Kim et al. 2009; Shah et al., 2009; Levey et al., 2011; Mahboubi et al., 2013). It therefore seems appropriate to review the main issues of concern. Seven main factors that have a significant effect on both the interpretation of results of individual studies and the extent to which the results of different studies can be meaningfully synthesised, were identified in this review. Each of these factors is described in the following section.

Inherent imprecision of PTA

As Schlauch and Carney (2012) emphasise, although PTA has been considered the “gold-standard” for assessing hearing threshold sensitivity (Shargorodsky et al. 2010), it is subject to variability due to calibration issues, test protocol, test-retest reliability, test environment, tester,

and participant factors (e.g., motivation). These factors become critical when attempting to identify incipient, or minimal, hearing loss (Schlauch and Carney, 2012), and where comparisons are made between data sets.

Influence of “pass-fail” criterion

In the clinical context, hearing thresholds better than 20 dB are generally treated as within “normal” limits (Lutman & Davis, 1994). However, the cut-off criterion (or “fence”) between “normal hearing” and “hearing loss”, has not been standardised among scientific investigators (Mehra et al., 2009; Shargorodsky et al., 2010). Many different criteria have been applied (some involving averaging of HTLs), generally without any clearly articulated justification. In estimating the prevalence of hearing loss within a population, the lower (i.e., stricter) the criterion adopted, the higher the reported prevalence (Mehra et al., 2009). A study by Lees et al. (1985), listed in Table 3.3, is a case in point. In this analysis, a very strict criterion was used (any HTL \geq 10 dB HL) with a resulting reported prevalence rate (40%) that far exceeds other estimates.

Reference HTL data

With respect to hearing loss criterion, it is also important to be mindful that “audiometric zero” is not an absolute, but must be inferred statistically from specific and adequate population data (Corso, 1963; citing Hirsch, 1952). Such population data is presented in ISO 7029 (ISO, 1984). General discrepancies between measured group thresholds and ISO 7029 data (ISO, 1984) have been reported by a number of investigators (Guest et al., 2012). If the ISO 7029 reference levels are not, in fact, typical of the general population, overestimation of prevalence will result, that is, the underestimation of “audiometric zero” will contribute to an assumption that experimental group data are intrinsically poorer than would otherwise be deduced. “Low

fence” estimates of hearing loss will also be additive with overly restrictive audiometric norms in this respect. In reviewing the results of audiometric surveys from various countries, Borchgrevink (2003) concluded that the median HTLs of any group of 18- to 20-year-olds are not 0 dB, but in the order of + 5 dB for most frequencies (0.25 – 8 kHz). This assertion is supported by recent Australian data (Williams et al., 2014). Schlauch and Carney (2011) noted that NHANES III (1988 – 1994) median HTLs are greater (i.e., worse) than 0 dB HL at each frequency, and suggested that this is not surprising, given that NHANES participants were not as stringently selected as those for studies contributing to ISO 7029. Reference databases that are carefully obtained and relevant to specific research target populations (e.g., adolescents and young adults) are currently lacking (ISO, 2013).

Baseline PTA

Given the uncertainties around reference audiometric data, NITS can really only be ascertained when pre-exposure (baseline) audiometric results are available. The majority of hearing surveys have not included baseline audiometry (Holgers & Petterson, 2005). Nevertheless, the terminology NITS (or NIPTS) has been used by some authors (e.g., Niskar et al., 2001). In the absence of baseline data, this is presumptive (Meinke & Dice, 2007) and may be misleading. It is also incorrect to assume that every individual starts with a pre-exposure 0 dB HL “baseline” even in the absence of other risk factors for hearing loss, as evidenced in ISO 7029 data.

Audiometric configuration

It has been clinically observed that subsequent to noise exposure, audiograms often show a frequency-specific hearing loss – typically in the 2 to 8 kHz region (Patuzzi, 1992) commonly referred to as a “noise notch”. The criterion used to identify a notch varies among investigators

(Mostafapour, 1998; Rabinowitz et al., 2006a), and determines the reported prevalence of its occurrence (Nondahl et al. 2009). Nondahl et al. (2009) systematically compared four notch “algorithms” (Coles et al., 2000; Dobie & Rabinowitz, 2002; McBride & Williams, 2001; Hoffman et al., 2006), observing poor agreement across criteria. Bilger (1976, p. 458) also cautioned that while group average notch data may be of interest, it must be ascertained whether a “typical” notch profile can also be systematically identified in noise-exposed individuals.

It is clear from the literature that, regardless of criterion used, not all individuals identified as having a noise notch report a positive history of noise exposure, and neither do all individuals reporting a positive history of noise exposure have a notch configuration (McBride & Williams, 2001; Nondahl et al., 2009; Osei-Lah & Yeoh, 2010). There is also evidence that “dips” at 4 or 6 kHz may have aetiological factors apart from noise exposure (e.g., genetic, viral infections, otitis media, skull trauma and ototoxic drugs; Sataloff, 1980; Haapaniemi, 1995). Klockhoff and Lyttkens (1982) presented 30 cases of children with a 4 kHz “dip”, none of whom had a history of noise exposure.

Reliance on “noise notches” occurring at 6 kHz is particularly problematic. Threshold elevation at 6 kHz may occur due to error in calibration reference values (Lutman & Qasam, 1998; McBride & Williams, 2001; Schlauch & Carney, 2011). Even small systematic errors such as these have a significant effect on the estimates of occurrence of notched audiograms. Schlauch and Carney (2011) re-analysed NHANES data, (using similar inclusion criteria to Niskar et al., 2001). They concluded that systematic threshold error at 6 kHz (for all age groups) and 8 kHz (for younger participants), were very likely to have influenced findings. Despite all these difficulties, a number of reporters on leisure-noise effects have used the noise notch as a “proxy” (Green, 2002) indicator of NIHL. Examples of some studies using the noise notch as a metric, and others critiquing the use of the notch, are listed in Table 3.6.

Table 3.6: Cross sectional surveys using audiometric configuration (notch) as indicator.

Authors	Participants	Assessment method	Findings
<i>Guild (1950)</i>	Unknown U.K.		“Abrupt” audiometric configurations not always associated with impulse noise exposure
<i>Hinchcliffe (1959)</i>	<i>N</i> = 100 U.K.	PTA; ENT exam; questionnaire re; risk factors	Poorer mid-high frequency HTLs in males – associated with small arms use
<i>Cozad (1974)</i>	<i>N</i> = 18,600 U.S. school students	PTA (no exposure data)	Cite audiometric configuration (hearing loss above 3000 Hz) as possible evidence of NIHL
<i>Axelsson (1981)</i>	<i>N</i> = 538 Swedish 17–20 yrs	PTA	15% showed some hearing loss. Refers to “dip” at 6 kHz – but no correlation with leisure-time activities
<i>Rytzner and Rytzner (1981)</i>	<i>N</i> = 14,391 Swedish 7, 10, 13 yrs	PTA screen; ENT follow up	Small occurrence (4 kHz “dip”) associated with exposure in approximately 200 cases
<i>Molvaer et al. (1983)</i>	<i>N</i> = 1474 Norwegian 20–80 yrs	PTA; ENT exam; questionnaire	6 kHz “dip” noted, even in youngest participants Assumed noise-related
<i>Haapenniemi (1995)</i>	<i>N</i> = 687 Finnish 6–15 yrs	PTA; ENT exam; questionnaire	8.3% occurrence Cites several possible factors
<i>Holmes et al. (1997)</i>	<i>N</i> = 342 U.S. 10–20 yrs	PTA screen; 6-item questionnaire	6kHz “dip” associated with firearms use
<i>Mostafapour (1998)</i>	<i>N</i> = 50 U.S. 18–30 yrs	PTA; speech discrimination test	Found only one case of “notch” at 6 kHz
<i>Niskar et al. (2001)</i>	<i>N</i> = 5249 U.S. 6–19 yrs	PTA (NHANES III data) (no exposure data)	12.5% estimated to have a “notch” (“NITS”)
<i>McBride and Williams (2001)</i>	<i>N</i> = 357 U.K. Electricity employees	Bekesy audiometry; questionnaire	49% determined to have a “notch” – no association with NIHL risk factors
<i>Rabinowitz et al. (2006a)</i>	<i>N</i> = 2526 U.S. 15–25 yrs	PTA (limited exposure data)	Almost 20% had “notch” – rate constant over 20 yr interval Likely related ‘at least in part’ to noise exposure

<i>Nondahl et al. (2009)</i>	<i>N</i> = 2395 U.S. 43–84 yrs	Compared algorithms using previous data (Beaver Dam study)	“Notches” noted in the absence of noise exposure history Poor agreement among four different algorithms
<i>Osei-Lah and Yeoh (2010)</i>	<i>N</i> = 149 U.K. outpatients 19–91 yrs	ENT outpatient assessment	39.6% exhibited “notches” not attributable to noise or other risk factors
<i>Schlauch and Carney (2011)</i>	<i>N</i> = 5089 U.S. 6–19 yrs	PTA (NHANES III data) compared test and re-test data and computer-simulated audiograms	Similar prevalence of “notches” in actual and simulated audiograms
<i>Jin et al. (2013)</i>	<i>N</i> = 698 U.S. marching band members and controls ≤ 25 yrs	PTA; OAE	Noted transitory behaviour of “notches” on multiple retests
<i>Twardella et al. (2013)</i>	<i>N</i> = 1843 German adolescents ~15–16 yrs	PTA	2.4% prevalence of “notches”

Apart from the scientific limitations of the use of the notch as a metric, it is also concerning that some papers confidently cited noise notch presence as evidence of NIHL, yet provided no substantiating data regarding participants’ actual leisure-noise exposure (e.g., Cozad et al., 1974; Niskar et al., 2001).

Explanatory variables

For meaningful estimates of NITS to be made, all possible risk factors for hearing loss must be taken into account (Engdahl et al., 2005). HTL data must be excluded from the analysis where factors other than noise exposure may contribute to HTL shift. Some extraneous factors can be observed at assessment (e.g., cerumen occlusion, middle ear dysfunction), while others must be identified through careful history taking (e.g., prenatal exposure to disease, ototoxic drug

exposure, family history, and head/ear trauma). There is also increasing evidence that other agents, such as tobacco (Ferrite & Santana, 2005) and solvents (Campo & Lataye, 2000), may represent significant risk factors for hearing loss.

Achieving the appropriate suite of exclusions is important, but challenging, as Schlauch and Carney (2012) demonstrate well. Obtaining an adequate case history is time consuming and subjective, as it relies on the recollection of the participant or informant. Further, a stringent set of exclusion criteria has the advantage of removing extraneous causes of variation, but introduces the disadvantage of decreased statistical power in particular strata (Lutman & Davis, 1994). There are insufficient test items in some data sets for strong exclusion criteria to be applied, which weakens the usefulness of the data in determining NIHL. NHANES III, for instance, did not include otoscopy or pure tone audiometric bone conduction testing in its protocol, a point that is clearly acknowledged by Niskar et al. (1998) and Shargorodsky et al. (2010). Tympanometry was included in the NHANES III protocol but Niskar et al. (1998) and Shargorodsky et al. (2010) apparently did not use the available results as an exclusion criterion in their analyses. The NHANES III data set also lacks information about the noise-exposure history of participants. Although the 2005 to 2006 NHANES protocol included a detailed questionnaire including probes on firearms use, occupational and non-occupational noise exposure, Shargorodsky et al. make only limited reference to this information in their commentary.

To demonstrate the importance of such analysis decisions, Schlauch and Carney (2012) re-analysed NHANES III and NHANES 2005 to 2006 data using the same criteria for hearing loss as Niskar et al. (1998) and Shargorodsky et al. (2010), but applying various exclusion criteria. They clearly demonstrated how criterion for hearing loss and exclusion criteria

interact to determine the estimated prevalence of hearing loss overall. Schlauch and Carney (2012) also highlighted the work of Henderson et al. (2011), who analysed the same data for teenage participants as Shargorodsky et al. (2010) but applied different exclusion criteria and a different criterion for hearing loss. Henderson et al. reported a lower prevalence of hearing loss than Shargorodsky et al. However, it is difficult to ascertain whether exclusions used, or the definition of hearing loss, contributed more to the difference in prevalence estimates (Schlauch and Carney, 2012).

It is also noteworthy that the analysis by Schlauch and Carney (2012) showed a higher percentage of 6- to 8-year-old participants met the criterion for hearing loss than 6- to 11-year-old children (18% versus 16.3%, respectively). Given that children as young as 6 to 8 years are unlikely to have any significant risk for NIHL, this finding casts further doubt on the assumption that the observed hearing losses in these young cohorts are attributable to noise exposure.

General comments

There are extensive data indicating that significant loud-noise exposure occurs in a range of leisure situations (in particular, when using firearms and attending rock-concerts, nightclubs and similar venues). However, there still appears to be insufficient consistent, empirical evidence to support the position that pure tone hearing loss, which is causally related to leisure-noise exposure, is either very widespread among young populations or is increasing over time. Much of the past research provides little insight into the lasting effects of leisure-noise exposure (e.g., experimental TTS studies, retrospective analyses of population data) and there is a dearth of more revealing studies (e.g., longitudinal studies). Some of the earlier literature provides insufficient detail for useful retrospective interpretation. The limitations of PTA as a

metric have not always been acknowledged. Case studies account for a large proportion of published material across a wide range of disciplines; however the scarcity of case studies in the leisure-noise literature (Luxon, 1998) is noteworthy. Although as a research methodology case studies are regarded as at the lower end of the evidence-based research pyramid, they can add depth of understanding to enquiries that large sample approaches do not provide and may help “close in” on real-life situations (Flyvbjerg, 2011). Perhaps most importantly, results from different studies using similar methods, or even using the same data, are not in good agreement.

Based on the evidence available for occupational noise exposure, some commentators have seemingly over-stated the likely longer-term effects of leisure-noise exposure. For example, Niskar et al. (2001) reported that children were found to have “moderate to profound NITS” and that “With continued harmful noise exposures, the threshold shift at 3, 4 or 6 kHz increases in severity...” (Niskar et al., 2001, p.40). Annex E of ISO 1999 (1990) indicates that the median threshold shift, even after four decades of very high intensity industrial noise exposure is moderate – not severe or profound – in degree, and the observed deterioration is gradual and non-linear, reaching an asymptote in time. For example, Table E.4 (ISO, 1990, p. 16) shows that the 0.5 fractile (i.e., median threshold value for the population) at 4000 Hz, after regular exposure to 100 dB ($L_{Aeq,8h}$) for ten years is 31 dB; for 20 years 36 dB; 30 years 39 dB and; 40 years 41 dB. This represents decreasing increments of 31, 5, 3 and finally, 2 dB per respective decade, in response to long and intense regular exposure. It is also noteworthy that the population data presented in ISO 1999 demonstrate that age-related threshold shift typically overlaps NITS later in life, as illustrated in Table B.1 of the standard (ISO, 1990).

Nevertheless, although the extent of the risk may have been overstated by some commentators, recent evidence (including “daily life” SPL measures) indicates that a proportion of young

people *are* exposed to noise doses sufficient to cause injury (Beach, 2013c). Beach et al. (2013c) recently estimated that ~ 15% may be affected. Although more conservative (lower) than some suggestions, this still represents a significant proportion of the population. The current authors agree with others (e.g., Smith, 2000; Schlauch & Carney, 2012) that public education is of continuing importance. However, it is equally important that the leisure-noise risk is not over-stated, at the potential cost of losing public credibility, and also diverting attention from the serious and ongoing problem of occupational noise. The additive effects of work, non-work and purely recreational noise sources must also be seriously considered (Williams, 2009). Despite the large volume of data collected in this field, no specific damage-risk criteria for leisure-noise exposure are currently available (Portnuff et al., 2013).

Implications for future research

Where PTA is used, it is imperative that protocols ensuring the highest level of precision are put into place (Macrae, 1998; Schlauch & Carney, 2007). Better quantification of the multiple sources of experimental uncertainty (e.g., measurement error, selection biases, confounding variables) is vital in designing studies that produce both meaningful and economical outcomes (Phillips, 2001). It also appears that the field would benefit greatly from the establishment of a consistent and scientifically justified approach to hearing loss criteria.

Previous commentators have suggested that paradigms other than conventional PTA should also be explored (e.g., SCENIHR, 2008). Suprathreshold tests (e.g., speech-in-noise assessments) and objective techniques (e.g., auditory brainstem response, cortical auditory evoked response, and MEG testing) may extend the evidence-base regarding subclinical

changes to the auditory system. Ideally, longitudinal studies would be implemented, with baseline measures collected in the pre-teen years when noise exposure is minimal. However, the current authors recognise the logistical challenges and high cost of this type of research. Thus, cross-sectional studies of clearly at risk groups (e.g., frequent nightclub attendees) may be useful. Amassing clinical details (including measures of hearing disability) of leisure-noise exposed individuals, with no other risk factors for hearing loss, may also be informative. The establishment of clinical databases to collate relevant information across localities, or even countries, could be considered.

The use of amplification potentially provides an additional risk factor for increased threshold shift, particularly in high-noise leisure environments. The current authors noted the lack of data concerning leisure-noise exposure and impacts for young hearing aid wearers (with early onset sensorineural hearing loss). This deficiency in knowledge is currently being addressed through an ongoing survey conducted by NAL. Lastly, in order to quantify the real individual and societal costs of leisure-noise exposure, more evidence regarding the actual hearing difficulties (i.e., disabling/handicapping effects) of leisure-noise-exposed individuals is greatly needed.

CONCLUSIONS

Some of the commentary in this field of research to date is arguably more speculative than evidence-based. Information provided to public health authorities, educationalists, the media, and the community at large, must be evidence-based and scientifically defensible. Scientists should be prepared to challenge over-stated, or over-paternalistic, public information. The real cost versus benefit of future leisure-noise research should be carefully considered, and the freedom of individuals to make personal choices about their recreational pursuits, based on

accurate scientific information, should be acknowledged in this process. At a global level, epidemiological and other health research directly consumes millions of dollars every year (Phillips, 2001). It is therefore imperative that future investigators do everything possible to ensure that unambiguous and meaningful conclusions can be reached in future leisure-noise research.

REFERENCES

- Ahmed, S., Y., Fallah, S., Garrido, B., et al. (2007). Use of portable audio devices by university students. *Canadian Acoustics*, 35, 35-48.
- Agrawal, Y., Platz, E. A., Niparko, J. K. (2008). Prevalence of hearing loss and differences by demographic characteristics among US adults: Data from the National Health and Nutrition Examination Survey, 1999–2004. *Arch Intern Med*, 168, 1522-1530.
- Axelsson, A., & F. Lindgren (1978). Temporary threshold shift after exposure to pop music. *Scand Audiol*, 7(3), 127-135.
- Axelsson, A., Jerson, T., Lindberg, U., et al. (1981). Early noise-induced hearing loss in teenage boys. *Scand Audiol*, 10(2), 91-96.
- Axelsson, A. (1991). Leisure noise exposure in adolescents and young adults. *J Sound Vib*, 151(3), 447-453.
- Axelsson, A., Rosenhall, U., Zachau, G. (1994). Hearing in 18-year-old Swedish males. *Scand Audiol*, 23(2), 129-134.
- Beach, E. F., Gilliver, M., Williams, W. (2013a). Leisure noise exposure: Participation trends, symptoms of hearing damage, and perception of risk. *Int J Audiol*, 52 Suppl 1, S20-S25.
- Beach, E., Gilliver, M., Williams, W. (2013b). The NOISE (Non-Occupational Incidents Situations and Events) database: A new research tool. *Ann Leisure Res*, 16(2), 149-159.
- Beach, E., & Nie, V. (2013). Noise levels in fitness classes are still too high: Evidence from 1997–98 and 2009–11. *Arch Environ Occup Health*, 69, 223-230. Available at: <http://dx.doi.org/10.1080/19338244.2013.771248>
- Beach, E., Williams, W., Gilliver, M. (2013c). Estimating young Australian adults' risk of hearing damage from selected leisure activities. *Ear Hear*, 34(1), 75-82.
- Biassoni, E. C., Serra, M. R., Richter, U., et al. (2005). Recreational noise exposure and its effects on the hearing of adolescents. Part II: Development of hearing disorders. *Int J Audiol*, 44(2), 74-85.
- Bienvenue, G. R., Michael, P. L., Violon-Singer, J. R. (1976). The effect of high level sound exposure on the loudness difference limen. *Am Ind Hyg Assoc J*, 37(11), 628-635.
- Bilger, R. C. (1976). The audiometric profile of noise-induced hearing loss. In: Henderson, D., Hamernik, R.P., Dosanjh, D.S., et al. (Eds). *Effects of Noise on Hearing*. New York, Raven Press.
- Bittel, S. N., Freeman, B. A., Kemker, B. E. (2008). Investigation of toy exposure in children. *Seminars in Hearing*, 29(1), 10-18.

- Bohlin, M. C., Erlandsson, S.I. (2007). Risk behaviour and noise exposure among adolescents. *Noise Health*, 9(36), 55-63.
- Borchgrevink, H. M. (1988). One-third of 18-year-old male conscripts show noise-induced hearing loss > 20 dB before start of military service – the incidence being doubled since 1981. Reflecting increased leisure noise? In: Berglund, U., Karlsson, J., Lindvall, T., (Eds). *Proceedings of the 5th international congress on noise as a public health problem*. Council for Building Research, August 21-28, 1988; Stockholm. 27-32.
- Borchgrevink, H. M. (2003). Does health promotion work in relation to noise? *Noise Health*, 5(18), 25-30.
- Brookhouser, P. E., Worthington, D. W., Kelly, W. J. (1992). Noise-induced hearing loss in children. *Laryngoscope*, 102(6), 645-655.
- Buffe, P., Cudennec, Y.F., Ben Azzouz, M., et al. (1986). [Survey on the harmfulness of listening to music with headphones. Article in French]. *Ann Otolaryngol Chir Cervicofac*, 103(5), 351-355.
- Campo, P., Lataye, R. (2000). Noise and solvent, alcohol and solvent: Two dangerous interactions on auditory function. *Noise Health*, 3(9), 49-57.
- Carter, N. L., Keen, K., Waugh, R. L., et al. (1978). *Hearing levels and oto-rhino-laryngological findings in students of the University of Sydney under age 21*. NAL Report No. 72. Canberra: National Acoustics Laboratories, Australian Government Publishing Service.
- Carter, N. L., Waugh, R. L., Keen, et al. (1982). Amplified music and young people's hearing. Review and report of Australian findings. *Med J Aust*, 2(3), 125-128.
- Carter, N. L., Murray, N., Khan, A., et al. (1984). A longitudinal study of recreational noise and young people's hearing. *Aust J Audiol*, 6(2), 45-53.
- Carter, L. (2011). Prevalence of hearing loss and its relationship to leisure sound exposure. Report to: Australian Government, Department of Health and Ageing, Canberra. Available at: http://www.health.gov.au/internet/main/publishing.nsf/Content/health-hear-prevalence_of_hearing_loss
- Centers for Disease Control and Prevention (2013). Available at: <http://www.cdc.gov/nchs/nhanes.htm>
- Chung, J. H., Des Roches, C. M., Meunier, J., et al. (2005). Evaluation of noise-induced hearing loss in young people using a web-based survey technique. *Pediatrics*, 115(4), 861-867.
- Clark, W. W. 1991. Noise exposure from leisure activities, a review. *J Acoust Soc Am*, 90(1), 175-181.
- Clark, W. W., Bohne, B. A. (1986). Temporary hearing losses following attendance at a rock concert. *J Acoust Soc Am*, Supp 1 (79) S48.
- Coles, R. R., Lutman, M. E., Buffin, J. T. (2000). Guidelines on the diagnosis of noise-induced hearing loss for medicolegal purposes. *Clin Otolaryngol*, 25, 264-273.
- Cone, B. K., Wake, M., Tobin, S., et al. (2010). Slight-mild sensorineural hearing loss in children, Audiometric: Clinical, and risk factor profiles. *Ear Hear*, 31(2), 202-212.
- Corso, J. F. (1963). Age and sex differences in pure-tone thresholds. Survey of hearing levels from 18 to 65 years. *Arch Otolaryngol*, 77, 385-405.
- Costa, O. A., Axelsson, A., Aniansson, G. (1988). Hearing loss at age 7, 10 and 13– an audiometric follow-up study. *Scand Audiol Suppl*, 30, 25-32.
- Cozad, R. L., Marston, L., Joseph, D. (1974). Some implications regarding high frequency hearing loss in school-age children. *J School Health*, 44, 92.
- Cranston, C. J., Brazile W. J., Sandfort, D. R., et al. (2013). Occupational and recreational noise exposure from indoor arena hockey games. *J Occup Environ Hyg*, 10(1), 11-16.
- Davis, A. C., Lovell, E. A., Smith, P. A., et al. (1998). The contribution of social noise to tinnitus in young people – A preliminary report. *Noise Health*, 1(1), 40-46.

- Dobie, R. A., Rabinowitz, P. M. (2002). Change in audiometric configuration helps to determine whether a standard threshold shift is work-related. *Spectrum* 19 Suppl 1, 17.
- Drake-Lee, A. B. (1992). Beyond music, auditory temporary threshold shift in rock musicians after a heavy metal concert. *J R Soc Med*, 85(10), 617-619.
- Emmerich, E., Richter, F., Hagner, H., et al. (2002). Effects of discotheque music on audiometric results and central acoustic evoked neuromagnetic responses. *Int Tinnitus J*, 8(1), 13-19.
- Engdahl, B., Tambs, K., Borchgrevink, H. M., et al. (2005). Screened and unscreened hearing threshold levels for the adult population: Results from the Nord-Trondelag Hearing Loss Study. *Int J Audiol*, 44(4), 213-230.
- Fearn, R. W. (1981). Hearing levels in school children aged 9-12 years and 13-16 years associated with exposure to amplified pop music and other noisy activities. *J Sound Vib*, 74, 151-153.
- Ferrite, S., & Santana, V. (2005). Joint effects of smoking, noise exposure and age on hearing loss. *Occup Med (Lond)*, 55(1), 48-53.
- Flamme, G. A., Stephenson, M. R., Deiters, K., et al. (2012). Typical noise exposure in daily life. *Int J Audiol*, 51 Suppl 1, S3-11.
- Flyvbjerg, B. (2006). Five misunderstandings about case-study research. *Qualitative Inquiry*, 12(2), 219-245.
- Green, J. (2002). Noise-induced hearing loss. *Pediatrics*, 109(5), 987-988.
- Guest, M., Boggess, M., Attia, J. (2012). Relative risk of elevated hearing threshold compared to ISO 1999 normative populations for Royal Australian Air Force male personnel. *Hear Res*, 285(1-2), 65-76.
- Guild, S. R. (1950). The progression of impaired hearing for high tones during childhood. *Laryngoscope*, 60(9), 885-911.
- Hanson, D. R., & Fearn, R. W. (1975). Hearing acuity in young people exposed to pop music and other noise. *Lancet*, 2(7927), 203-205.
- Haapaniemi, J. (1995). The 6 kHz acoustic dip in school-aged children in Finland. *Eur Arch Otorhinolaryngol*, 252(7), 391-394.
- Hellström, P., Axelsson, A., Costa, O. (1998). Temporary threshold shift induced by music. *Scand Audiol*, 48, 87-94.
- Henderson, D., Subramaniam, M., Boettcher, F. A. (1993). Individual susceptibility to noise-induced hearing loss, an old topic revisited. *Ear Hear*, 14(3), 152-168.
- Henderson, E., Testa, M. A., Hartnick, C. (2011). Prevalence of noise-induced hearing-threshold shifts and hearing loss among US Youths. *Pediatrics*, 127(1), e39-46.
- Hètu, R., & Fortin, M. (1995). Potential risk of hearing damage associated with exposure to highly amplified music. *J Am Acad Audiol*, 6, 378-386.
- Hidecker, M. J. C. (2008). Noise-induced hearing loss in school-age children: What do we know? *Seminars in Hearing*, 29(1), 19-28.
- Hill, A. B. (1965). The environment and disease, association or causation? *Proc R Soc Med*, 58, 295-300.
- Hinchcliffe, R. (1959). The threshold of hearing in a random sample rural population. *Acta Oto-laryngol*, 50, 411-422.
- Hirsch, I. J. (1952). *The Measurement of Hearing*. New York, McGraw-Hill Book Company, Inc.
- Hoffman, H. J., Ko, C.-W., Themann, C.L., et al. (2006). Reducing noise-induced hearing loss (NIHL) to achieve U.S. Healthy People 2010 goals. *Am J Epidemiol*, 163 Suppl S122. Available at: http://aje.oxfordjournals.org/content/163/suppl_11.toc

- Hoffman, H. J., Dobie, R. A., Ko, C-W., et al. (2010). Americans hear as well or better today compared with 40 years ago: Hearing threshold levels in the unscreened adult population of the United States, 1959–1962 and 1999–2004. *Ear Hear*, 31(6), 725-734.
- Holgers, K., & Pettersson B. (2005). Noise exposure and subjective hearing symptoms among school children in Sweden. *Noise Health*, 7, 27-37.
- Holmes, A. E., Kaplan, H. S., Phillips, R. M., et al. (1997). Screening for hearing loss in adolescents. *Lang Speech Hear Serv Sch*, 28, 70-76.
- International Standards Organization (ISO) (1984). *ISO 7029: Acoustics – Threshold of hearing by air conduction as a function of age and sex for otologically normal persons*. Geneva, International Organization for Standardization.
- ISO (1990). *ISO 1999: Acoustics – Determination of occupational noise exposure and estimation of noise-induced hearing impairment*. Geneva, International Organization for Standardization.
- ISO (2013). *ISO 1999 Third edition: Acoustics – Estimation of noise-induced hearing loss*. Switzerland, International Organization for Standardization.
- Jin, S.-H., Nelson, P.B., Schlauch, R.S., et al. (2013). Hearing conservation program for marching band members: A risk for noise-induced hearing loss? *Am J Audiol*, 22, 26-39.
- Jorge Junior, J. J. (1993). *Evaluation of the hearing thresholds of young listeners of amplified music*. Doctoral dissertation, [in Portuguese]. Department of Otolaryngology, University of São Paulo, Brazil.
- Kardous, C., & Morata, T. (2010). Occupational and recreational noise exposures at stock car racing circuits: An exploratory survey of three professional race tracks. *Noise Control Eng*, 58(1), 54-61.
- Keppler, H., Dhooge, I., Maes, L., et al., (2010). Short-term auditory effects of listening to an MP3 player. *Arch Otolaryngol Head Neck Surg*, 136(6), 538-548.
- Kim, M. G., Hong, S. M., Shim, H. J., et al. (2009). Hearing threshold of Korean adolescents associated with the use of personal music players. *Yonsei Med J*, 50(6), 771-776.
- Klockhoff, I., & Lyttkens, L. (1982). Hearing defects of noise trauma type with lack of noise exposure. *Scand Audiol*, 11(4), 257-260.
- Kujawa, S. G., & Liberman, M. C. (2009). Adding insult to injury: Cochlear nerve degeneration after "temporary" noise-induced hearing loss. *J Neurosci*, 29(45), 14077-14085.
- Kumar, A., Ameenudin, S., Sangamanatha, A.V. (2012). Temporal and speech processing skills in normal hearing individuals exposed to occupational noise. *Noise Health*, 14(58), 100-105.
- Lee, P. C., Senders, C. W., Gantz, et al. (1985). Transient sensorineural hearing loss after overuse of portable headphone cassette radios. *Otolaryngol Head Neck Surg*, 93(5), 622-625.
- Lees, R. E., Roberts, J. H., Wald, Z. (1985). Noise induced hearing loss and leisure activities of young people, a pilot study. *Can J Public Health*, 76(3), 171-173.
- Le Prell, C. G., Hensley, B. N., Campbell, K. C., et al. (2011). Evidence of hearing loss in a 'normally-hearing' college-student population. *Int J Audiol*, 50 Suppl 1, S21-31.
- Levey, S., Levey, T., Fligor, B. J. (2011). Noise exposure estimates of urban MP3 player users. *J Speech Lang Hear Res*, 54(1), 263-277.
- Lindgren, F., Axelsson, A. (1983). Temporary threshold shift after exposure to noise and music of equal energy. *Ear Hear*, 4(4), 197-201.
- Lutman, M. E., & Davis, A. C. (1994). The distribution of hearing threshold levels in the general population aged 18-30 years. *Audiology*, 33(6), 327-350.

- Lutman, M. E., & Qasam, H.Y. (1998). A source for notches at 6.0 kHz. In: Prasher, D., Luxon, L. (Eds). *Advances in Noise Research, Biological Effects of Noise Vol I* (pp. 170-176). London, Whurr Publishers Ltd.
- Luxon, L. (1998). Toys and games, poorly recognised hearing hazards? *BMJ*, 316, 1473-1480.
- Maassen, M., Babisch, W., Bachmann, K. D., et al. (2001). Ear damage caused by leisure noise. *Noise Health*, 4(13), 1-16.
- McBride, D., & Williams, S. (2001). Characteristics of the audiometric notch as a clinical sign of noise exposure. *Scand Audiol*, 30(2), 106-111.
- Macrae, J. (1988). Improvement of Clinical Monitoring Audiometry. *Aust J Audiol*, 10(1), 12-15.
- McCombe, A. W., Binnington, J., Davis, A., et al. (1995). Hearing loss and motorcyclists. *J Laryngol Otol*, 109(7), 599-604.
- McFadden, S. L., Henderson, D., Shen, Y. H. (1997). Low-frequency 'conditioning' provides long-term protection from noise-induced threshold shifts in chinchillas. *Hear Res*, 103(1-2), 142-150.
- McMillan, P. M., & Kileny, P. R. (1994). Hearing loss from a bicycle horn. *J Am Acad Audiol*, 5(1), 7-9.
- Mahboubi, H., Oliaei, S., Badran, K. W., et al. (2013). Systematic assessment of noise amplitude generated by toys intended for young children. *Otolaryngol Head Neck Surg*, 148(6), 1043-1047.
- Makary, C. A., Shin, J., Kujawa, S. G., et al. (2011). Age-related primary cochlear neuronal degeneration in human temporal bones. *J Assoc Res Otolaryngol*, 12(6), 711-717.
- Martinez-Wbaldo, M. d. C., Soto-Vazquez, C., Ferre-Calacich, I., et al. (2009). Sensorineural hearing loss in high school teenagers in Mexico City and its relationship with recreational noise. *Cad. Saúde Pública. Rio de Janeiro*, 25(12), 2553-2561.
- Mazelova, J., Valvoda, J., Popelar, J., et al. (2001). The influence of single exposure to amplified music on hearing of young listeners. In: Henderson, D., Prasher, D., Kopke, R., et al. (Eds). *Noise Induced Hearing Loss, Basic Mechanisms, Prevention and Control*. London, nRn Publications, 365-375.
- Mehra, S., Eavey, R. D., Keamy, D. G., Jr. (2009). The epidemiology of hearing impairment in the United States, newborns, children, and adolescents. *Otolaryngol Head Neck Surg*, 140(4), 461-472.
- Meinke, D. K., Dice, N. (2007). Comparison of audiometric screening criteria for the identification of noise-induced hearing loss in adolescents. *Am J Audiol*, 16(2), S190-202.
- Meyer-Bisch, C. (1996). Epidemiological evaluation of hearing damage related to strongly amplified music (personal cassette players, discotheques, rock concerts) – high-definition audiometric survey on 1364 subjects. *Audiology*, 35(3), 121-142.
- Miller, J. D. (1974). Effects of noise on people. *J Acoust Soc Am*, 56(3), 729-764.
- Mills, J. H. (1975). Noise and children: A review of literature. *J Acoust Soc Am*, 58, 767-779.
- Mirbod, S. M., Inaba, R., Yoshida, H., et al. (1992). Noise exposure level while operating electronic arcade games as a leisure time activity. *Ind Health*, 30(2), 65-76.
- Molvaer, O. I., Vardal, L., Gundersen, et al. (1983). Hearing acuity in a Norwegian standard population. *Scand Audiol*, 12(4), 229-236.
- Morata, T. C. (2007). Young people, their noise and music exposures and the risk of hearing loss. *Int J Audiol*, 46(3), 111-112.
- Mostafapour, S. P., Lahargoue, K., Gates, G. A. (1998). Noise-induced hearing loss in young adults, the role of personal listening devices and other sources of leisure noise. *Laryngoscope*, 108(12), 1832-1839.

- Medical Research Council Institute of Hearing Research (MRC) (1986). Damage to hearing arising from leisure noise. *Br J Audiol*, 20, 157-164.
- Nassar, G. (2001). The human temporary threshold shift after exposure to 60 minutes' noise in an aerobics class. *Br J Audiol*, 35(1), 99-101.
- Neitzel, R., Seixas, N., Olson, J., et al. (2004). Nonoccupational noise, Exposures associated with routine activities. *J Acoust Soc Am*, 115(1), 237-245.
- Nicotera, T., Henderson, D., Hu, B. H., et al. (2001). Noise exposure and mechanisms of hair cell death. In: Henderson, D., Prasher, D., Kopke, R., et al. (Eds). *Noise Induced Hearing Loss. Basic Mechanisms, Prevention and Control*. London, nRn Publications, p. 99-117.
- Niskar, A. S., Kieszak, S. M., Holmes, A., et al. (1998). Prevalence of hearing loss among children 6 to 19 years of age: The third National Health and Nutrition Examination Survey. *JAMA*, 279(14), 1071-1075.
- Niskar, A. S., Kieszak, S. M., Holmes, A. E., et al. (2001). Estimated prevalence of noise-induced hearing threshold shifts among children 6 to 19 years of age: the Third National Health and Nutrition Examination Survey, 1988–1994, United States. *Pediatrics*, 108(1), 40-43.
- Niu, X., Canlon, B. (2002). Protective mechanisms of sound conditioning. *Adv Otorhinolaryngol*, 59, 96-105.
- Nondahl, D. M., Shi, X., Cruickshanks, K. J., et al. (2009). Notched audiograms and noise exposure history in older adults. *Ear Hear*, 30(6), 696-703.
- Okamoto, H., Teismann, H., Kakigi, R., et al. (2011). Broadened population-level frequency tuning in human auditory cortex of portable music player users. *PLoS One*, 6(3), e17022.
- Op de Beeck, K., Schacht, J., Van Camp, G. (2011). Apoptosis in acquired and genetic hearing impairment. The programmed death of the hair cell. *Hear Res*, 281(1-2), 18-27.
- Opperman, D. A., Reifman, W., Schlauch, R., et al. (2006). Incidence of spontaneous hearing threshold shifts during modern concert performances. *Otolaryngol Head Neck Surg*, 134(4), 667-673.
- Osei-Lah, V., & Yeoh, L. H. (2010). High frequency audiometric notch, an outpatient clinic survey. *Int J Audiol*, 49(2), 95-98.
- Pääkkönen R., Jokitulppo, J., Heinijoki, V.M., et al. (2003). Noise exposure of spectators at Finnish air shows. *Applied Acoustics*, 64(2), 121-127.
- Park, M. Y. (2003). Assessment of potential noise-induced hearing loss with commercial karaoke noise. *Int J Ind Ergon*, 31, 375-385.
- Patuzzi, R. (1992). Effects of noise on auditory nerve responses. In: Dancer, A., Henderson, D., Salvi, R.J., et al. (Eds). *Noise-Induced Hearing Loss*. St Louis, MO, Mosby Year Book, Inc.
- Peng, J-H., Tao, Z-Z., Huang, Z-W. (2007). Risk of damage to hearing from personal listening devices in young adults. *J Otolaryngol*, 36(3), 181-185.
- Persson, B. O., Svedberg, A., Gothe, C.J. (1993). Longitudinal changes in hearing ability among Swedish conscripts. *Scand Audiol*, 22, 141-143.
- Petrescu, N. (2008). Loud music listening. *Mcgill J Med*, 11, 169-176.
- Phillips, C. V. (2001). The economics of 'more research is needed'. *Int J Epidemiol*, 30, 771-776.
- Phillips, C. V., & Goodman, K. J. (2004). The missed lessons of Sir Austin Bradford Hill. *Epidemiol Perspect Innov*, 1(1), 3.
- Portnuff, C.D.F., Fligor, B. J., Arehart, K. H. (2013). Self-report and long-term field measures of MP3 player use, how accurate is self-report? *Int J Audiol*, 52 Suppl 1, S33-40.

- Punch, J. L., Elfenbein, J. L., James, R. R. (2011). Targeting hearing health messages for users of personal listening devices. *Am J Audiol*, 20(1), 69-82.
- Quaranta, A., Sallustio, V., Quaranta, N. (2001). Noise induced hearing loss, Summary and perspectives. In: Henderson, D., Prasher, D., Kopke, R., et al. (Eds). *Noise Induced Hearing Loss. Basic Mechanisms, Prevention and Control*. London, nRn Publications, 539-557.
- Rabinowitz, P. M., Galusha, D., Slade, M. D., et al. (2006a). Audiogram notches in noise-exposed workers. *Ear Hear*, 27, 742-750.
- Rabinowitz, P. M., Slade, M. D., Galusha, D., et al. (2006b). Trends in the prevalence of hearing loss among young adults entering an industrial workforce 1985 to 2004. *Ear Hear*, 27(4), 369-375.
- Ramsey, K. L., Simmons, F. B. (1993). High-powered automobile stereos. *Otolaryngol Head Neck Surg*, 109(1), 108-110.
- Reddell, R. C. & C. P. Lebo (1972). Ototraumatic effects of hard rock music. *Calif Med*, 116(1), 1-4.
- Rintelmann, W. F., & Borus, J. F. (1968). Noise-induced hearing loss and rock and roll music. *Arch Otolaryngol*, 88(4), 377-385.
- Roche, A. F., Siervogel, R. M., Himes, J. H., et al. (1977). *Longitudinal study of human hearing: Its relationship to noise and other factors. I. Design of Five Year Study; Data from First Year*. Ohio, USA, Wright-Patterson Air Force Base. AMRL-TR-76-110 (ADA040168).
- Roche, A. F., Himes, J. H., Siervogel, R. M. (1979). *Longitudinal study of human hearing: Its relationship to noise and other factors II. Results from the First Three Years*. Ohio, USA, Wright-Patterson Air Force Base.
- Roche, A. F., Chumlea, W. M. C., Siervogel, R. M., et al. (1982). *Longitudinal study of human hearing: Its relationship to noise and other factors. Raw data: Hearing threshold and noise exposure. Final report* (pp. 1-66). Yellow Springs, Ohio: Aerospace Medical Research Laboratory, Aerospace Medical Division, Air Force Systems Command, Wright Patterson Airforce Base, Ohio. AFAMRL-TR-83-057 (AD-A148-112).
- Rose, A. S., Ebert, C. S., Jr., Prazma, J., et al. (2008). Noise exposure levels in stock car auto racing. *Ear Nose Throat J*, 87(12), 689-692.
- Rytzner, B., Rytzner, C. (1981). Schoolchildren and noise. The 4 kHz dip – tone screening in 14391 schoolchildren. *Scand Audiol*, 10(4), 213-216.
- Sadhra, S., Jackson, C. A., Ryder, et al. (2002). Noise exposure and hearing loss among student employees working in university entertainment venues. *Ann Occup Hyg*, 46(5), 455-463.
- Sataloff, R. T. (1980). The 4000-Hz audiometric dip. *Ear Nose Throat J*, 59 24-32.
- Scientific Committee on Emerging and Newly Identified Health Risks (SCENIHR) (2008). Potential health risks of exposure to noise from personal music players and mobile phones including a music player function. *Scientific Committee on Emerging and Newly Identified Health Risks, European Commission*, 1-81.
- Schlauch, R. S., & Carney, E. (2007). A multinomial model for identifying significant pure-tone threshold shifts. *J Speech Lang Hear Res*, 50(6), 1391-1403.
- Schlauch, R. S., & Carney, E. (2011). Are false-positive rates leading to an overestimation of noise-induced hearing loss? *J Speech Lang Hear Res*, 54(2), 679-692.
- Schlauch, R. S., & Carney, E. (2012). The challenge of detecting minimal hearing loss in audiometric surveys. *Am J Audiol*, 21, 106-119.
- Schmidt, J. M., Verschuure, J., Brocaar, M. P. (1994). Hearing loss in students at a conservatory. *Audiology*, 33(4), 185-194.

- Serra, M. R., Biassoni, E. C., Richter, U., et al. (2005). Recreational noise exposure and its effects on the hearing of adolescents. Part I: An interdisciplinary long-term study. *Int J Audiol*, 44(2), 65-73.
- Shah, S., Gopal, B., Reis, J., et al. (2009). Hear today, gone tomorrow: An assessment of portable entertainment player use and hearing acuity in a community sample. *J Am Board Fam Med*, 22(1), 17-23.
- Shargorodsky, J., Curhan, S. G., Curhan, G. C., et al. (2010). Change in prevalence of hearing loss in US adolescents. *JAMA*, 304(7), 772-778.
- Smith, P. A., Davis, A., Ferguson, M., et al. (2000). The prevalence and type of social noise exposure in young adults in England. *Noise Health*, 2(6), 41-56.
- Strasser, H., Irle, H., Scholz, R. (1999). Physiological cost of energy-equivalent exposures to white noise, industrial noise, heavy metal music, and classical music. *Noise Control Eng*, 47 (5), 187-192.
- Strasser, H., Irle, H., Legler, R. (2003). Temporary hearing threshold shifts and restitution after energy-equivalent exposures to industrial noise and classical music. *Noise Health*, 5(20), 75-84.
- Strauss, P., Quante, M., Strahl, M., et al. (1977). [Is hearing of the students damaged by environmental noise in their leisure time? (author's transl)] [article in German]. *Laryngol Rhinol Otol (Stuttg)*, 56(10), 868-871.
- Swanson, S. J., Dengerink, H. A., Kondrick, P., et al. (1987). The influence of subjective factors on temporary threshold shifts after exposure to music and noise of equal energy. *Ear Hear*, 8(5), 288-291.
- Tam, R., Sharma, M., Williams, W., et al. (2013). The effect of listening to MP3 music on otoacoustic emissions. *Aust NZ J Audiol*, 33(2) 125-134.
- Tambs, K., Hoffman, H. J., Borchgrevink, H. M., et al. (2003). Hearing loss induced by noise, ear infections, and head injuries: Results from the Nord-Trondelag Hearing Loss Study. *Int J Audiol*, 42(2), 89-105.
- Tharpe, A. M., Sladen, D.P. (2008). Causation of permanent unilateral and mild bilateral hearing loss in children. *Trends Amplif*, 12(1), 17-25.
- Tin, L. L., Lim, O. P. (2000). A study on the effects of discotheque noise on the hearing of young patrons. *Asia Pac J Public Health*, 12(1), 37-40.
- Torre III, P., & Howell, J. C. (2008). Noise levels during aerobics and the potential effects on distortion product otoacoustic emissions. *J Commun Disord*, 41(6), 501-511.
- Turunen-Rise, I., Flottorp, G., Tvete, O. (1991). Personal cassette players ('Walkman'). Do they cause noise-induced hearing loss? *Scand Audiol*, 20(4), 239-244.
- Twardella, D., Perez Alvarez, C., Steffens, T., et al. (2011). [Hearing loss in adolescents due to leisure noise. The OHRKAN study] [article in German, with English abstract]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz*, 54(8), 965-971.
- Twardella, D., C. Perez-Alvarez, Steffans, T., et al. (2013). The prevalence of audiometric notches in adolescents in Germany: The Ohrkan-study. *Noise Health*, 15(67), 412-419.
- Vittitow, M., Windmill, I. M., Yates, J. W., et al. (1994). Effect of simultaneous exercise and noise exposure (music) on hearing. *J Am Acad Audiol*, 5(5), 343-348.
- Wang, Y., Hirose, K., Liberman, M. C. (2002). Dynamics of noise-induced cellular injury and repair in the mouse cochlea. *J Assoc Res Otolaryngol*, 3(3), 248-268.
- Ward, W. D. (1976). A comparison of the effects of continuous, intermittent and impulse noise. In: Henderson, D., Hamernik, R.P., Dosanjh, D.S., et al. (Eds). *Effect of Noise on Hearing*. New York, Raven Press, 407-420.
- West, P. D., Evans, E. F. (1990). Early detection of hearing damage in young listeners resulting from exposure to amplified music. *Br J Audiol*, 24(2), 89-103.

- Williams, W. (2009). Noise exposure profiles. Acoustics 2009, Adelaide, Australia, Australian Acoustical Society.
- Williams, W. (2011). Barriers to Occupational Noise Management. *The reduction of noise exposure in the workplace*. Saarbrücken, LAP LAMBERT Academic Publishing.
- Williams, W., Carter, L., Seeto, M. (2014). Hearing threshold levels for a population of 11 to 35 year old Australian females and males. *Int J Audiol*, 53, 289-293.
- World Health Organization (WHO) (1980). *Environmental Health Criteria 12. Noise*. Geneva, World Health Organization.
- Yaremchuk, K., Dickson, L., Burk, K., et al. (1997). Noise level analysis of commercially available toys. *Int J Pediatr Otorhinolaryngol*, 41(2), 187-197.
- Yassi, A., Pollock, N., Tran, N., et al. (1993). Risks to hearing from a rock concert. *Can Fam Physician*, 39, 1045-1050.
- Ylikoski, J., Xing-Qun, L., Virkkala, J., et al. (2001). Molecular mechanisms of the death of auditory hair cells by noise. In: Henderson, D., Hamernik, R.P, Dosanjh, D.S., et al. (Eds). *Noise Induced Hearing Loss, Basic Mechanisms, Prevention and Control*. London, nRn Publications, 89-98.
- Zhao, F., Manchaiah, V. K., French, D., et al. (2010). Music exposure and hearing disorders: An overview. *Int J Audiol*, 49(1), 54-64.
- Zocoli, A. M., Morata, T.C., Marques, J.M., et al. (2009). Brazilian young adults and noise, attitudes, habits, and audiological characteristics. *Int J Audiol*, 48(10), 692-699.